

## DESCRIPTION

## BICYCLOAMIDE DERIVATIVE

## TECHNICAL FIELD

[0001]

5           The present invention relates to bicycloamide derivatives and pharmaceutically acceptable salts thereof that inhibit dipeptidylpeptidase IV (DPP-IV) and are useful in the prevention and/or treatment of type II diabetes and other diseases that involve DPP-IV.

## 10 BACKGROUND ART

[0002]

          Dipeptidylpeptidase IV (EC3.4.14.5, referred to as "DPP-IV" or "CD26," hereinafter) is a serine protease that specifically hydrolyzes polypeptides having proline or alanine  
15 at position 2 on the C-terminal side of these amino acid residues, cleaving dipeptides Xaa-Pro or Xaa-Ala from the N-terminus of the polypeptides (Xaa may be any amino acid).

[0003]

          One biological function of DPP-IV is the inactivation of  
20 glucagon-like peptide 1 (GLP-1) by cleaving the N-terminal His-Ala dipeptide of GLP-1 (Non-Patent Document 1). The GLP-1 inactivated by DPP-IV is thought to act as an antagonist on GLP-1 receptors, further decreasing the physiological activity of GLP-1 (Non-Patent Document 2). GLP-1, a peptide hormone  
25 secreted from endocrine L-cells found primarily in intestinal

epithelium, is known to act on  $\beta$ -cells of the pancreatic Langerhans' islets in a glucose-dependent manner to promote the insulin secretion, thus decreasing the blood glucose level (Non-Patent Documents 3 and 4). Having an ability to promote  
5 insulin biosynthesis and  $\beta$ -cell growth, GLP-1 is an essential factor for the maintenance of  $\beta$ -cells (Non-Patent Documents 5 and 6). It has been reported that GLP-1 also acts to promote glucose utilization by peripheral tissue and, when intraventricularly administered, decreases food intake and  
10 motility of GI tract (Non-Patent Documents 7 through 10).  
[0004]

A DDP-IV inhibitor is believed to increase the GLP-1 activity by suppressing the decomposition of innate GLP-1. The increased GLP-1 activity stimulates insulin secretion and  
15 improves glucose metabolism. For this reason, DPP-IV inhibitors are expected to be useful in the prevention and/or treatment of diabetes, in particular type II diabetes (Non-Patent Documents 11 and 12). The compounds are expected to be also effective in the prevention and/or treatment of other  
20 diseases that are caused or worsened by decreased glucose metabolism (for example, diabetic complications, hyperinsulinemia, hyperglycemia, abnormal lipid metabolism and obesity).  
[0005]

25 The roles of DPP-IV in a living body other than the

inactivation of GLP-1 and how the enzyme is involved in the onset of various diseases have been described in many reports as described below.

[0006]

5 (a) DPP-IV inhibitors and their antibodies prevent the invasion of HIV into cells. Expression of CD26 is reduced in T-cells derived from patients infected with HIV-1 (Non-Patent Document 13). HIV-1 Tat protein binds to DPP-IV (Non-Patent Document 14).

10 [0007]

(b) DPP-IV is involved in immune responses. DPP-IV inhibitors and their antibodies suppress the growth of T-cells stimulated by antigens (Non-Patent Document 15). T-cells stimulated by antigens express an increased level of DPP-IV (Non-Patent Document 16). DPP-IV is involved in the cytokine production and other functions of T-cells (Non-Patent Document 17). DPP-IV binds to adenosine deaminase (ADA) on the T-cell surface (Non-Patent Document 18).

[0008]

20 (c) Expression of DPP-IV is increased in the skin fibroblasts of patients with rheumatoid arthritis, psoriasis, and lichen planus (Non-Patent Document 19).

[0009]

(d) High DPP-IV activity is observed in patients with  
25 benign prostatic hypertrophy and in the homogenate of the

prostatic tissue (Non-Patent Document 20). DPP-IV in the lung endothelium acts as an adhesive molecule for lung-metastatic breast cancer and prostatic cancer in rats (Non-Patent Document 21).

5 [0010]

(e) The DPP-IV defective variant of F344 rats has lower blood pressure than the wild-type F344 rats. DPP-IV interacts with a protein that plays a crucial role in sodium reabsorption by the kidney (Patent Documents 1 and 2).

10 [0011]

(f) The inhibition of DPP-IV activity offers an effective approach to the prevention and/or treatment of myelosuppressive diseases, while DPP-IV-activating agents are expected to serve as drugs to increase the white blood cell count and/or treat infectious diseases (Patent Document 3).

15 [0012]

These observations indicate that DPP-IV inhibitors can be useful in the prevention and/or treatment of diabetes (in particular, type II diabetes) and/or diseases other than diabetic complications that involve DPP-IV. For example, DPP-IV inhibitors are expected to be useful in the prevention and/or treatment of AIDS following infection with HIV, rejection following organ/tissue transplantation, multiple sclerosis, rheumatoid arthritis, inflammation, allergies, osteoporosis, psoriasis and lichen planus, benign prostatic

20  
25

hypertrophy, lung metastasis of breast and prostatic cancers, hypertension and infectious diseases. DPP-IV inhibitors are also expected to be used to facilitate diuresis, decrease myelosuppression and increase white blood cell count.

5 [0013]

Among existing DPP-IV inhibitors are pyrrolidine derivatives described in Patent Documents 4 through 11, heterocyclic derivatives described in Patent Documents 12 and 13, and  $\beta$ -amino acid derivatives described in Patent Documents  
10 14 and 15.

[0014]

Patent Document 16, a US patent, discloses a single bicycle[2.2.2]octane derivative that inhibits DPP-IV activity. This compound, however, is completely different from the  
15 compounds of the present invention in its structure and mechanism for DPP-IV inhibition. Patent Document 17 mentions a bicycle derivative structurally similar to the compounds of the present invention. However, there is no description in this literature concerning any of the compounds of the present  
20 invention, nor have any examples been presented of the compounds.

[0015]

None of the previously described DPP-IV inhibitors are practical enough in terms of DPP-IV inhibitory activity,  
25 selectivity for DPP-IV, stability, toxicity and biological

kinetics. Thus, a constant need exists for effective DDP-IV inhibitors.

[Non-Patent Document 1] American Journal of Physiology, Vol. 271 (1996): ppE458-E464.

5 [Non-Patent Document 2] European Journal of Pharmacology, Vol. 318 (1996): pp429-435

[Non-Patent Document 3] European Journal Clinical Investigation, Vol. 22 (1992): p154

[Non-Patent Document 4] Lancet, Vol. 2 (1987): p1300

10 [Non-Patent Document 5] Endocrinology, Vol. 42 (1992): p856

[Non-Patent Document 6] Diabetologia, Vol. 42 (1999):p 856

[Non-Patent Document 7] Endocrinology, Vol. 135 (1994): p2070

[Non-Patent Document 8] Diabetologia, Vol. 37 (1994): p1163

[Non-Patent Document 9] Digestion, Vol. 54 (1993): p392

15 [Non-Patent Document 10] Dig. Dis. Sci., Vol. 43 (1998): p1113

[Non-Patent Document 11] Diabetes, Vol. 47 (1998): pp1663-1670

[Non-Patent Document 12] Diabetologia, Vol. 42 (1999):  
20 pp1324-1331

[Non-Patent Document 13] Journal of Immunology, Vol. 149 (1992): p3037

[Non-Patent Document 14] Journal of Immunology, Vol. 150 (1993): p2544

25 [Non-Patent Document 15] Biological Chemistry (1991): p305

- [Non-Patent Document 16]        Scandinavian Journal of Immunology, Vol. 33 (1991): p737
- [Non-Patent Document 17]        Scandinavian Journal of Immunology, Vol. 29 (1989): p127
- 5    [Non-Patent Document 18]        Science, Vol. 261 (1993): p466
- [Non-Patent Document 19]        Journal of Cellular Physiology, Vol. 151 (1992): p378
- [Non-Patent Document 20]        European Journal of Clinical Chemistry and Clinical Biochemistry, Vol. 30 (1992): p333
- 10   [Non-Patent Document 21]        Journal of Cellular Physiology, Vol. 121 (1993): p1423
- [Patent Document 1]        WO 03/015775 Pamphlet
- [Patent Document 2]        WO 03/017936 Pamphlet
- [Patent Document 3]        WO 03/080633 Pamphlet
- 15   [Patent Document 4]        WO 95/15309 Pamphlet
- [Patent Document 5]        WO 98/19998 Pamphlet
- [Patent Document 6]        WO 00/34241 Pamphlet
- [Patent Document 7]        WO 02/14271 Pamphlet
- [Patent Document 8]        WO 02/30890 Pamphlet
- 20   [Patent Document 9]        WO 02/38541 Pamphlet
- [Patent Document 10]        WO 03/002553 Pamphlet
- [Patent Document 11]        US 02/0193390 Publication
- [Patent Document 12]        WO 02/062764 Pamphlet
- [Patent Document 13]        WO 03/004496 Pamphlet
- 25   [Patent Document 14]        WO 03/000180 Pamphlet

[Patent Document 15] WO 03/004498 Pamphlet

[Patent Document 16] US 02/0193390 Publication

[Patent Document 17] WO 02/38541 Pamphlet

#### DISCLOSURE OF THE INVENTION

#### 5 PROBLEMS TO BE SOLVED BY THE INVENTION

[0016]

It is an object of the present invention to provide a novel compound that has high DPP-IV inhibitory activity, as well as pharmaceutically acceptable salts thereof. It is  
10 another object of the present invention to provide a pharmaceutical composition containing the novel compound that has high DPP-IV inhibitory activity or a pharmaceutically acceptable salt thereof. It is still another object of the present invention to provide a prophylactic and/or therapeutic  
15 agent for diabetes and associated complications, as well as a prophylactic and/or therapeutic agent for diseases involving DPP-IV.

#### MEANS TO SOLVE THE PROBLEMS

[0017]

20 According to the present invention, there are provided a novel bicycloamide derivative that has high DPP-IV inhibitory activity, and pharmaceutically acceptable salts thereof. Also provided is a pharmaceutical composition containing the novel bicycloamide derivative that has high DPP-IV inhibitory  
25 activity, or a pharmaceutically acceptable salt thereof.




Further provided are a prophylactic and/or therapeutic agent for diabetes and associated complications, and a prophylactic and/or therapeutic agent for diseases involving DPP-IV.

[0018]

5 Thus, the present invention concerns the following:

1) A bicycloamide derivative represented by the following general formula (1):

[0019]




10 [0020]

[wherein  $R^1$  and  $R^2$  may or may not be identical to one another and are each independently a hydrogen atom, substituted or unsubstituted  $C_1$  to  $C_6$  alkyl group, substituted or unsubstituted  $C_3$  to  $C_6$  cycloalkyl group, substituted or unsubstituted arylmethyl group, substituted or unsubstituted arylethyl group, substituted or unsubstituted aromatic hydrocarbon group, substituted or unsubstituted aromatic heterocyclic ring, substituted or unsubstituted aliphatic heterocyclic ring or  $NR^3R^4$  (wherein  $R^3$  and  $R^4$  may or may not be identical to one another and are each independently a hydrogen atom, substituted or unsubstituted  $C_1$  to  $C_6$  alkyl group, substituted or unsubstituted  $C_3$  to  $C_6$  cycloalkyl group,

substituted or unsubstituted arylmethyl group, substituted or  
 unsubstituted aromatic hydrocarbon group, substituted or  
 unsubstituted aromatic heterocyclic ring or substituted or  
 unsubstituted aliphatic heterocyclic ring, or  $R^3$  and  $R^4$  may  
 5 together form a ring structure.), or  $R^1$  and  $R^2$  may together  
 form a ring structure; X is  $CH_2$ , CHF,  $CF_2$ , CHOH, S or O; and n  
 is 1, 2 or 3.],  
 or a pharmaceutically acceptable salt thereof.

2) The bicycloamide derivative as set forth in 1) above,  
 10 represented by the following general formula (2):

[0021]




[0022]

[wherein  $R^5$  is a substituted or unsubstituted  $C_1$  to  $C_6$  alkyl  
 15 group, substituted or unsubstituted  $C_3$  to  $C_6$  cycloalkyl group,  
 substituted or unsubstituted arylmethyl group, substituted or  
 unsubstituted arylethyl group, substituted or unsubstituted  
 aromatic hydrocarbon group, substituted or unsubstituted  
 aromatic heterocyclic ring, substituted or unsubstituted  
 20 aliphatic heterocyclic ring or  $NR^3R^4$  (wherein  $R^3$  and  $R^4$  may or  
 may not be identical to one another and are each independently  
 a hydrogen atom, substituted or unsubstituted  $C_1$  to  $C_6$  alkyl

group, substituted or unsubstituted C<sub>3</sub> to C<sub>6</sub> cycloalkyl group, substituted or unsubstituted arylmethyl group, substituted or unsubstituted aromatic hydrocarbon group, substituted or unsubstituted aromatic heterocyclic ring or substituted or unsubstituted aliphatic heterocyclic ring, or R<sup>3</sup> and R<sup>4</sup> may together form a ring structure.); X is CH<sub>2</sub>, CHF, CF<sub>2</sub>, CHOH, S or O; and n is 1, 2 or 3.],  
or a pharmaceutically acceptable salt thereof.

3) A bicycloamide derivative as set forth in 1) above,  
represented by the following general formula (3):

[0023]




[0024]

[wherein R<sup>7</sup> and R<sup>8</sup> may or may not be identical to one another and are each independently a substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl group, substituted or unsubstituted C<sub>3</sub> to C<sub>6</sub> cycloalkyl group, substituted or unsubstituted arylmethyl group, substituted or unsubstituted arylethyl group, substituted or unsubstituted aromatic hydrocarbon group, substituted or unsubstituted aromatic heterocyclic ring, substituted or unsubstituted aliphatic heterocyclic ring or NR<sup>3</sup>R<sup>4</sup> (wherein R<sup>3</sup> and R<sup>4</sup> may or may not be identical to one

another and are each independently a hydrogen atom,  
 substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl group, substituted  
 or unsubstituted C<sub>3</sub> to C<sub>6</sub> cycloalkyl group, substituted or  
 unsubstituted arylmethyl group, substituted or unsubstituted  
 5 aromatic hydrocarbon group, substituted or unsubstituted  
 aromatic heterocyclic ring or substituted or unsubstituted  
 aliphatic heterocyclic ring, or R<sup>3</sup> and R<sup>4</sup> may together form a  
 ring structure.), or R<sup>7</sup> and R<sup>8</sup> may together form a ring  
 structure; X is CH<sub>2</sub>, CHF, CF<sub>2</sub>, CHOH, S or O; and n is 1, 2 or  
 10 3.],  
 or a pharmaceutically acceptable salt thereof.

4) An intermediate in the production of the bicycloamide  
 derivative of 1) above, represented by the following formula  
 (4):



[0026]

[wherein R<sup>1</sup> and R<sup>2</sup> may or may not be identical to one another  
 and are each independently a hydrogen atom, substituted or  
 20 unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl group, substituted or  
 unsubstituted C<sub>3</sub> to C<sub>6</sub> cycloalkyl group, substituted or  
 unsubstituted arylmethyl group, substituted or unsubstituted

arylethyl group, substituted or unsubstituted aromatic hydrocarbon group, substituted or unsubstituted aromatic heterocyclic ring, substituted or unsubstituted aliphatic heterocyclic ring or  $\text{NR}^4\text{R}^5$  (wherein  $\text{R}^4$  and  $\text{R}^5$  may or may not be identical to one another and are each independently a hydrogen atom, substituted or unsubstituted  $\text{C}_1$  to  $\text{C}_6$  alkyl group, substituted or unsubstituted  $\text{C}_3$  to  $\text{C}_6$  cycloalkyl group, substituted or unsubstituted arylmethyl group, substituted or unsubstituted aromatic hydrocarbon group, substituted or unsubstituted aromatic heterocyclic ring or substituted or unsubstituted aliphatic heterocyclic ring, or  $\text{R}^4$  and  $\text{R}^5$  may together form a ring structure.), or  $\text{R}^1$  and  $\text{R}^2$  may together form a ring structure; X is  $\text{CH}_2$ ,  $\text{CHF}$ ,  $\text{CF}_2$ ,  $\text{CHOH}$ , S or O; n is 1, 2 or 3; and  $\text{P}^1$  is an amino-protecting group].

5) A pharmaceutical product, containing as an active ingredient the bicycloamide derivative as set forth in 1) above or a pharmaceutically acceptable salt thereof.

6) A DPP-IV inhibitor, containing as an active ingredient the bicycloamide derivative as set forth in 1) above or a pharmaceutically acceptable salt thereof.

7) A therapeutic agent for treating diseases involving DPP-IV, containing as an active ingredient the bicycloamide derivative as set forth in 1) above or a pharmaceutically acceptable salt thereof.

[0027]

As used herein, the term "substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl group" refers to a C<sub>1</sub> to C<sub>6</sub> alkyl group (such as methyl group, cyclopropylmethyl group, ethyl group, propyl group, 1-methylethyl group, 1-methylpropyl group, 2-methylpropyl group, 1-ethylpropyl group, 2-ethylpropyl group, butyl group, t-butyl group and hexyl group) that may contain 1 to 5 substituents selected from halogen atom, hydroxy group, cyano group, C<sub>1</sub> to C<sub>6</sub> alkoxy group, substituted or unsubstituted aryloxy group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonyl group, C<sub>1</sub> to C<sub>6</sub> alcoxycarbonyl group, C<sub>1</sub> to C<sub>6</sub> alkylthio group, amino group, mono- or di-substituted C<sub>1</sub> to C<sub>6</sub> alkylamino group, 4- to 9-membered cyclic amino group that may contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alcoxycarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkylsulfonylamino group, substituted or unsubstituted arylsulfonylamino group and other substituents.

[0028]

As used herein, the term "substituted or unsubstituted C<sub>3</sub> to C<sub>6</sub> cycloalkyl group" refers to a C<sub>3</sub> to C<sub>6</sub> cycloalkyl group (such as cyclopropyl group, cyclobutyl group, cyclopentyl group and cyclohexyl group) that may contain 1 to 5 substituents selected from halogen atom, hydroxy group, cyano group, C<sub>1</sub> to C<sub>6</sub> alkoxy group, substituted or unsubstituted aryloxy group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonyl group, C<sub>1</sub> to C<sub>6</sub> alcoxycarbonyl group, C<sub>1</sub> to C<sub>6</sub> alkylthio group, amino group,

mono- or di-substituted C<sub>1</sub> to C<sub>6</sub> alkylamino group, 4- to 9-membered cyclic amino group that may contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkoxy carbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkylsulfonylamino group, substituted or unsubstituted arylsulfonylamino group and other substituents.

[0029]

As used herein, the term "substituted or unsubstituted arylmethyl group" refers to an arylmethyl group (such as phenylmethyl group, naphthylmethyl group, pyridylmethyl group, quinolylmethyl group and indolylmethyl group) that may contain 1 to 5 substituents selected from halogen atom, substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl group, hydroxy group, cyano group, nitro group, substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkoxy group, substituted or unsubstituted aryloxy group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonyl group, C<sub>1</sub> to C<sub>6</sub> alkoxy carbonyl group, C<sub>1</sub> to C<sub>6</sub> alkylthio group, amino group, mono- or di-substituted C<sub>1</sub> to C<sub>6</sub> alkylamino group, substituted or unsubstituted arylamino group, 4- to 9-membered cyclic amino group that may contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkoxy carbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkylsulfonylamino group, substituted or unsubstituted arylsulfonylamino group and other substituents.

[0030]

As used herein, the term "substituted or unsubstituted

arylethyl group" refers to an arylethyl group (such as 1-phenethyl group, 2-phenethyl group, 1-naphthylethyl group and 2-naphthylethyl group) that may contain 1 to 5 substituents selected from halogen atom, substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl group, hydroxy group, cyano group, nitro group, substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkoxy group, substituted or unsubstituted aryloxy group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonyl group, C<sub>1</sub> to C<sub>6</sub> alkoxycarbonyl group, C<sub>1</sub> to C<sub>6</sub> alkylthio group, amino group, mono- or di-substituted C<sub>1</sub> to C<sub>6</sub> alkylamino group, substituted or unsubstituted arylamino group, 4- to 9-membered cyclic amino group that may contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkoxycarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkylsulfonylamino group, substituted or unsubstituted arylsulfonylamino group and other substituents.

[0031]

As used herein, the term "substituted or unsubstituted aromatic hydrocarbon group" refers to an aromatic hydrocarbon group (such as benzene ring, naphthalene ring and anthracene ring) that may contain 1 to 5 substituents selected from halogen atom, hydroxy group, cyano group, nitro group, C<sub>1</sub> to C<sub>6</sub> alkoxy group, C<sub>1</sub> to C<sub>6</sub> alkylthio group, amino group, mono- or di-substituted C<sub>1</sub> to C<sub>6</sub> alkylamino group, 4- to 9-membered cyclic amino group that may contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub>



alkylsulfonylamino group, substituted or unsubstituted  
arylsulfonylamino group and other substituents.

[0032]

As used herein, the term "substituted or unsubstituted  
5 aromatic heterocyclic ring" refers to an aromatic heterocyclic  
ring (e.g., 5- or 6-membered aromatic monocyclic heterocyclic  
ring or 9- or 10-membered fused aromatic heterocyclic ring,  
such as pyridine ring, pyrimidine ring, pyridazine ring,  
triazine ring, quinoline ring, naphthyridine ring, quinazoline  
10 ring, acridine ring, pyrrole ring, furan ring, thiophene ring,  
imidazole ring, pyrazole ring, oxazole ring, isoxazole ring,  
thiazole ring, indole ring, benzofuran ring, benzothiazole  
ring, benzimidazole ring and benzoxazole ring. The  
heterocyclic ring contains 1 to 3 hetero atoms selected from  
15 nitrogen atom, oxygen atom and sulfur atom.) that may contain  
1 to 5 substituents selected from halogen atom, hydroxy group,  
cyano group, nitro group, C<sub>1</sub> to C<sub>6</sub> alkoxy group, C<sub>1</sub> to C<sub>6</sub>  
alkylthio group, amino group, mono- or di-substituted C<sub>1</sub> to C<sub>6</sub>  
alkylamino group, 4- to 9-membered cyclic amino group that may  
20 contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub>  
alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkylsulfonylamino group,  
substituted or unsubstituted arylsulfonylamino group and other  
substituents.

[0033]

25 As used herein, the term "substituted or unsubstituted

aliphatic heterocyclic ring" refers to an aliphatic heterocyclic ring (e.g., 4- to 7-membered aliphatic monocyclic heterocyclic ring or 9- or 10-membered fused aliphatic heterocyclic ring, such as azetidine ring, pyrrolidine ring, tetrahydrofuran ring, piperidine ring, morpholine ring and piperazine ring. The heterocyclic ring contains 1 to 3 hetero atoms selected from nitrogen atom, oxygen atom and sulfur atom.) that may contain 1 to 5 substituents selected from halogen atom, substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl group, hydroxy group, cyano group, substituted or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkoxy group, C<sub>1</sub> to C<sub>6</sub> alkylthio group, amino group, mono- or di-substituted C<sub>1</sub> to C<sub>6</sub> alkylamino group, 4- to 9-membered cyclic amino group that may contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkoxy carbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkylsulfonylamino group, substituted or unsubstituted arylsulfonylamino group and other substituents.

[0034]

As used herein, the term "substituted or unsubstituted alkoxy group" refers to a C<sub>1</sub> to C<sub>6</sub> alkoxy group (such as methoxy group, ethoxy group, butoxy group and hexyloxy group) that may contain 1 to 5 substituents selected from halogen atom, hydroxy group, cyano group, C<sub>1</sub> to C<sub>6</sub> alkoxy group, C<sub>1</sub> to C<sub>6</sub> alkylthio group, amino group, mono- or di-substituted C<sub>1</sub> to C<sub>6</sub> alkylamino group, 4- to 9-membered cyclic amino group that

may contain 1 to 3 hetero atoms, formylamino group, C<sub>1</sub> to C<sub>6</sub> alkylcarbonylamino group, C<sub>1</sub> to C<sub>6</sub> alkylsulfonylamino group, substituted or unsubstituted arylsulfonylamino group and other substituents. The term "amino-protecting group" as used  
5 herein refers to such substituents as t-butoxycarbonyl group, benzyloxycarbonyl group, allyloxycarbonyl group, methoxycarbonyl group, ethoxycarbonyl group, 2,2,2-trichloroethoxycarbonyl group, trifluoroacetyl group, acetyl group, benzyl group and 2,4,6-trimethoxybenzyl group. As used  
10 herein, the term "a ring that R<sup>1</sup> and R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup>, or R<sup>7</sup> and R<sup>8</sup> together form" refers to an aliphatic heterocyclic ring (e.g., 4- to 7-membered aliphatic monocyclic heterocyclic ring or 9- or 10-membered fused aliphatic heterocyclic ring, such as azetidine ring, pyrrolidine ring, piperidine ring, morpholine  
15 ring and piperazine ring. The heterocyclic ring contains 1 to 3 hetero atoms selected from nitrogen atom, oxygen atom and sulfur atom.), a benzo-analogue of aliphatic heterocyclic rings (e.g., 4- to 7-membered aliphatic monocyclic heterocyclic ring or 9- or 10-membered fused aliphatic  
20 heterocyclic ring, such as azetidine ring, pyrrolidine ring, piperidine ring, morpholine ring and piperazine ring. The heterocyclic ring contains 1 to 3 hetero atoms selected from nitrogen atom, oxygen atom and sulfur atom.), imidazole ring or benzimidazole ring. As used herein, the term "halogen  
25 atom" refers to fluorine atom, chlorine atom, bromine atom or

iodine atom.

[0035]

Among preferred examples of the compound of the present invention are (2S,4S)-1-[[N-(4-carbamoylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile and (2S)-1-[[N-(4-carbamoylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile.

#### ADVANTAGE OF THE INVENTION

[0036]

10 The present invention provides novel DPP-IV inhibitors that are useful not only in the prevention and/or treatment of diabetes and associated complications, but also in the prevention and/or treatment of other diseases involving DPP-IV.

#### BRIEF DESCRIPTION OF THE DRAWINGS

15 [0037]

Fig. 1 is a graph showing the effect of Compound 1 on the plasma glucose level in normal mice, as determined in the oral glucose tolerance test. Each plot is given as the average of five examples  $\pm$  standard deviation (T-test with  $P < 0.05$  vs control).

#### BEST MODE FOR CARRYING OUT THE INVENTION

[0038]

When the compounds of the present invention form pharmaceutically acceptable salts, they may form salts with inorganic acids, such as hydrochloric acid, hydrobromic acid,

sulfuric acid, nitric acid and phosphoric acid; organic acids, such as acetic acid, maleic acid, fumaric acid, succinic acid, lactic acid, malic acid, tartaric acid, citric acid, methanesulfonic acid, p-toluenesulfonic acid, benzenesulfonic acid, salicylic acid, stearic acid, palmitic acid and trifluoroacetic acid; metals, such as sodium, potassium, calcium, magnesium, aluminum and zinc; ammoniums, such as ammonium and tetramethylammonium; organic amines, such as morpholine and piperidine; and amino acids, such as glycine, lysine, arginine, phenylalanine, and proline.

[0039]

The compounds of the present invention represented by the general formula (1) or salts thereof may contain a single or two or more chiral centers and thus have multiple optical isomers resulting from these chiral centers. Any of these optical isomers and diastereomers are encompassed by the present invention, as are any mixtures thereof in an arbitrary mixing ratio, including racemic mixtures. When the compounds of the present invention represented by the general formula (1) or salts thereof contain a double bond, they may have Z- or E-configuration and any of the mixtures of these compounds in an arbitrary mixing ratio are also encompassed by the present invention. Some of the compounds of the present invention represented by the general formula (1) or salts thereof may have tautomers or rotational isomers, all of which

isomers are encompassed by the present invention, as are any of the mixtures thereof in an arbitrary mixing ratio.

[0040]

The compounds of the present invention represented by the  
5 general formula (1) or salts thereof include intramolecular salts, addition products, solvates, and hydrates thereof.

[0041]

The compounds of the present invention represented by the general formula (1) or salts thereof may be used as a  
10 pharmaceutical composition either individually or in conjunction with one or more pharmaceutically acceptable auxiliary agents: They may be formulated with pharmaceutically acceptable carriers or excipients (such as starch, lactose, calcium phosphate, and calcium carbonate), lubricants (such as  
15 magnesium stearate, calcium stearate talc, and stearic acid), binders (such as starch, crystalline cellulose, carboxy methyl cellulose, gum arabic, polyvinyl pyrrolidone, and alginic acid), disintegrating agents (such as talc and carboxy methyl cellulose calcium) or diluents (such as saline, aqueous  
20 solutions of glucose, mannitol or lactose). Using ordinary techniques, the compounds of the present invention represented by the general formula (1) or salts thereof may be formulated into tablets, capsules, granules, powders, subtle granules, ampoules, or injections for oral or parenteral administration.

25 The compounds of the present invention represented by the

general formula (1) or salts thereof are generally administered to humans and other mammals at a dose of 0.0001 to 1000mg/kg/day while the dose may vary depending on the type of the compound or salt, route of administration, and the age, body weight, and symptoms of the subjects. The compounds of the present invention or salts thereof may be administered in a single daily dose or multiple doses per day.

[0042]

When necessary, the compounds of the present invention represented by the general formula (1) or salts thereof may be used in conjunction with one or more diabetic therapeutic agents other than DPP-IV inhibitors. Among such diabetic therapeutic agents for use with the compounds of the present invention or salts thereof are insulin and its derivatives, GLP-1 and its derivatives, and other oral diabetic therapeutic agents. Examples of the oral diabetic therapeutic agents include sulfonyl urea diabetic therapeutic agents, non-sulfonylurea insulin secretagogues, biguanide diabetic therapeutic agents,  $\alpha$ -glycosidase inhibitors, glucagon antagonists, GLP-1 agonists, PPAR agonists,  $\beta$ 3 agonists, SGLT inhibitors, PKC inhibitors, glucagon synthase kinase 3 (GSK-3) inhibitors, protein tyrosine phosphatase 1B (PTP-1B) inhibitors, potassium channel openers, insulin sensitizers, glucose uptake modulators, compounds modifying lipid metabolism, and appetite suppressors.

[0043]

Examples of GLP-1 and its derivatives include betatropin and NN-2211. Examples of sulfonylurea diabetic therapeutic agents include tolbutamide, glibenclamide, gliclazide, glimepiride, and glipizide. Examples of non-sulfonylurea insulin secretagogues include nateglinide, repaglinide, mitiglinide, and JTT-608. Examples of biguanide diabetic therapeutic agents include metformin. Examples of  $\alpha$ -glycosidase inhibitors include voglibose and miglitol. Examples of PPAR agonists include troglitazone, rosiglitazone, pioglitazone, ciglitazone, KRP-297 (MK-767), isaglitazone, GI-262570, and JTT-501. Examples of  $\beta$ 3 agonists include AJ-9677, YM-178, and N-5984.

[0044]

The compounds (1) of the present invention can be produced by various synthetic techniques. The compounds (1) of the present invention can be isolated or purified by common separation means (such as extraction, recrystallization, distillation, and chromatography). The compounds may be obtained in the form of various salts by using common techniques or similar techniques (such as neutralization).

[0045]


Representative processes for producing the compounds of the present invention and salts thereof will now be described.

[0046]



Process A

[0047]



[0048]

5 Step 1 (Process A)


In this step, a haloacetic acid derivative of the general formula (6) (where Y<sup>1</sup> is Cl or Br, and X is as defined above.) is reacted with a bicycloamine derivative of the general formula (5) (where R<sup>1</sup>, R<sup>2</sup> and n are as defined above.) to  
10 obtain a bicycloamide derivative of claim 1 (where R<sup>1</sup>, n and X are as defined above.). The reaction is carried out in the presence or absence of a base. Examples of the base for use in this reaction may include an inorganic base, such as sodium hydroxide, potassium hydroxide, sodium bicarbonate, potassium bicarbonate, sodium carbonate, potassium carbonate and cesium  
15 carbonate, or an organic base, such as triethylamine, diisopropyl ethylamine, N,N,N,N-tetramethyl ethylenediamine, diazabicyclo[5.4.0]-7-undecene, diazabicyclo[4.3.0]-5-nonene, phosphazene base and pentaisopropylguanidine. Examples of the  
20 catalyst for use in this reaction may include a phase transfer catalyst or an inorganic salt, such as tetrabutyl ammonium

bromide, tetrabutyl ammonium iodide, benzyl triethyl ammonium bromide, lithium bromide, lithium iodide, sodium iodide, potassium bromide, potassium iodide, cesium bromide and cesium iodide. The solvent for use in the reaction may be an inert solvent that does not affect the reaction, including, for example, acetone, ethanol, toluene, acetonitrile, tetrahydrofuran, dioxane, ethylether, t-butyl methyl ether, dimethoxy ethane, ethyl acetate, dichloro methane, N,N-dimethyl formamide, dimethyl sulfoxide and N-methyl-2-pyrrolidone. This reaction proceeds smoothly at 0 to 150°C.

[0049]

Process B

[0050]



[0051]

Step 1 (Process B)

In this step, a haloacetic acid derivative of the general

formula (6) (where X and Y<sup>1</sup> are as defined above.) is reacted with a bicycloamine derivative of the general formula (7) (where P<sup>2</sup> is a protective group for a carboxyl group, and n is as defined above.) to obtain a bicycloamide derivative of the general formula (8) as set forth in claim 1 (where P<sup>2</sup>, n, and X are as defined above.). The reaction is carried out in the presence or absence of a base. Examples of the base for use in this reaction may include an inorganic base, such as sodium hydroxide, potassium hydroxide, sodium bicarbonate, potassium bicarbonate, sodium carbonate, potassium carbonate and cesium carbonate, or an organic base, such as triethylamine, diisopropyl ethylamine, N,N,N,N-tetramethyl ethylenediamine, diazabicyclo[5.4.0]-7-undecene, diazabicyclo[4.3.0]-5-nonene, phosphazene base and pentaisopropylguanidine. Examples of the catalyst for use in the reaction may include a phase transfer catalyst or an inorganic salt, such as tetrabutyl ammonium bromide, tetrabutyl ammonium iodide, benzyl triethyl ammonium bromide, lithium bromide, lithium iodide, sodium iodide, potassium bromide, potassium iodide, cesium bromide and cesium iodide. The solvent for use in the reaction may be an inert solvent that does not affect the reaction, including, for example, as acetone, ethanol, toluene, acetonitrile, tetrahydrofuran, dioxane, ethylether, t-butyl methyl ether, dimethoxy ethane, ethyl acetate, dichloro methane, N,N-dimethyl formamide, dimethyl sulfoxide and N-methyl-2-

pyrrolidone. This reaction proceeds smoothly at 0 to 150°C.

[0052]

#### Step 2 (Process B)

In this step, the secondary amino group of the  
5 bicycloamide derivative of the general formula (8) (where  $P^2$ , n  
and X are as defined above.) is protected to give a  
bicycloamide derivative of the general formula (9) as set  
forth in claim 1 (where  $P^1$  is a protective group for amino  
group, and  $P^2$ , n and x are as defined above.). The protective  
10 group  $P^1$  for the secondary amine group may be t-butoxycarbonyl  
group, benzyloxycarbonyl group or trifluoroacetyl group. The  
protective groups can be introduced by known techniques. For  
example, when  $P^1$  is t-butoxycarbonyl group, it can be readily  
introduced by reacting di-t-butylidicarbonate with the  
15 bicycloamide derivative of the general formula (8) (where  $P^2$ , n  
and X are as defined above.) in the presence or absence of  
triethylamine or N,N-dimethylaminopyridine. When  $P^1$  is  
benzyloxycarbonyl group, it can be readily introduced by  
reacting benzyloxycarbonyl chloride with the bicycloamide  
20 derivative of the general formula (8) (where  $P^2$ , n and X are as  
defined above.) in the presence of triethylamine, diisopropyl  
ethylamine or potassium carbonate. When  $P^1$  is trifluoroacetyl  
group, it can be readily introduced by reacting  
trifluoroacetic acid anhydride with the bicycloamide  
25 derivative of the general formula (8) (where  $P^2$ , n and X are as

defined above.) in the presence of triethylamine or 4-dimethylaminopyridine.

[0053]

#### Step 3 (Process B)

5           In this step, the  $P^2$  group that protects the carboxyl group of the bicycloamide derivative of the general formula (9) (where  $P^2$ ,  $P^1$ ,  $n$  and  $X$  are as defined above.) is removed to give a bicycloamide derivative of the general formula (10) as set forth in claim 1 (where  $P^1$ ,  $n$  and  $X$  are as defined above.).

10    $P^2$  can be removed by known techniques. When  $P^2$  is t-butyl group, it can be readily removed by using trifluoroacetic acid or a solution of hydrogen chloride/dioxane. When  $P^2$  is benzyl group, it can be readily removed by using palladium carbon in combination with hydrogen or ammonium formate. When  $P^2$  is

15   tetrahydropyranyl group, it can be readily removed by using acetic acid, p-toluenesulfonic acid or hydrochloric acid.

[0054]

#### Step 4 (Process B)

          In this step, the bicycloamide derivative of the general

20   formula (10) (where  $P^1$ ,  $n$  and  $X$  are as defined above.) and an amine derivative of the formula  $R^1R^2NH$  (where  $R^1$  and  $R^2$  are as defined above) are reacted in the presence of a condensation agent for amidation to give a bicycloamide derivative of the general formula (11) as set forth in claim 4 (where  $R^1$ ,  $R^2$ ,  $P^1$ ,

25    $n$  and  $X$  are as defined above.). Examples of the condensation

agents used in this step include dicyclohexylcarbodiimide (DCC), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), dimethylimidazolinium chloride (DMC), ethyl chloroformate, isobutyl chloroformate and pivaloyl chloride. These agents may be added in the form of solid, liquid or a solution in a proper solvent. Examples of the base for use in the condensation reaction may include an alkali carbonate, such as sodium bicarbonate and potassium carbonate, and a tertiary amine, such as triethylamine, diisopropyl ethylamine, N-methylmorpholine, diazabicyclo[5.4.0]-7-undecene, pyridine, 4-dimethylaminopyridine and 1,8-bis(dimethylamino)naphthalene. The solvent for use in the condensation reaction may be an inert solvent that does not affect the reaction, including, for example, N,N-dimethylformamide, N,N-dimethylacetamide, dimethyl sulfoxide, acetonitrile, tetrahydrofuran, dioxane, ethyl ether, dimethoxyethane, ethyl acetate, toluene and dichloromethane. This condensation reaction proceeds smoothly at -20 to 150°C.

[0055]

#### Step 5 (Process B)

In this step, the bicycloamide derivative of the general formula (10) (where  $P^1$ , n and X are as defined above.) is converted to a bicyclic derivative of the general formula (12)

[where W is a reaction residue (such as halogen atoms, and

halides, imidazolides and active esters of carboxylic acids, such as 1-imidazolyl group, 4-nitrophenoxy group, pentafluorophenoxy group, imidoyloxy succinate group and 1-benzotriazolyloxy group (or 1-benzotriazolyl 3-oxide group),  $P^1$ ,  
5 n and X are as described above.]. This step can be readily carried out by known techniques. When W is imidoyloxy succinate group, the bicycloamide derivative of the general formula (10) (where  $P^1$ , n and X are as defined above.) is reacted with N-hydroxysuccinic acid in the presence of a  
10 condensation agent to give the desired product. When W is benzotriazolyloxy group (or 1-benzotriazolyl 3-oxide group), the bicycloamide derivative of the general formula (10) (where  $P^1$ , n and X are as defined above.) is reacted with 1-hydroxybenzotriazole in the presence of a condensation agent  
15 to give the desired product. Examples of the condensation agent for use in this step include dicyclohexylcarbodiimide (DCC), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), dimethylimidazolinium chloride (DMC), ethyl chloroformate, isobutyl chloroformate and pivaloyl  
20 chloride. These agents may be added in the form of solid, liquid or a solution in a proper solvent. When it is desired to use a base in the condensation reaction, examples of the base include an alkali carbonate, such as sodium bicarbonate and potassium carbonate, and a tertiary amine, such as  
25 triethylamine, diisopropyl ethylamine, N-methylmorpholine,

diazabicyclo[5.4.0]-7-undecene, pyridine, 4-

dimethylaminopyridine and 1,8-bis(dimethylamino)naphthalene.

The solvent for use in the condensation reaction may be an inert solvent that does not affect the condensation reaction,

5 including, for example, N,N-dimethylformamide, N,N-

dimethylacetamide, dimethylsulfoxide, acetonitrile,

tetrahydrofuran, dioxane, ethyl ether, dimethoxyethane, ethyl acetate, toluene and dichloromethane. This condensation

reaction proceeds smoothly at -20 to 150°C. The resulting  
10 bicyclic derivative of the general formula (12) (where W, P<sup>1</sup>, n and X are as described above.) may be used in the subsequent step after purification or as the unpurified crude product.

[0056]

#### Step 6 (Process B)

15 In this step, the bicycloamide derivative of the general formula (12) (where W, P<sup>1</sup>, n and X are as described above.) is reacted with an amine derivative of the formula R<sup>1</sup>R<sup>2</sup>NH (where R<sup>1</sup> and R<sup>2</sup> are as defined above.) to give a bicycloamide

derivative of the general formula (11) as set forth in claim 4  
20 (where R<sup>1</sup>, R<sup>2</sup>, P<sup>1</sup>, n and X are as defined above.). When a base is used in the reaction, the base may be an inorganic salt, such as sodium hydroxide, potassium hydroxide, sodium

bicarbonate, potassium bicarbonate, sodium carbonate, potassium carbonate and cesium carbonate, or an organic base,  
25 such as triethylamine, diisopropyl ethylamine, N,N,N,N-



tetramethylethylenediamine, diazabicyclo[5.4.0]-7-undecene, diazabicyclo[4.3.0]-5-nonene, phosphazine base and pentaisopropylguanidine. The solvent for use in the reaction may be an inert solvent that does not affect the reaction, such as toluene, acetonitrile, tetrahydrofuran, dioxane, ethylether, t-butylmethylether, dimethoxyethane, ethyl acetate, dichloromethane, N,N-dimethylformamide, dimethylsulfoxide and N-methyl-2-pyrrolidone. This reaction proceeds smoothly at - 30 to 150°C.

10 [0057]


#### Step 7 (Process B)

In this step, the P<sup>1</sup> group that protects the secondary amino group in the bicycloamide derivative of the general formula (11) (where R<sup>1</sup>, R<sup>2</sup>, P<sup>1</sup>, n and X are as defined above.) is removed to give a bicycloamide derivative of the general formula (1) as set forth in claim 1 (where R<sup>1</sup>, R<sup>2</sup>, n and X are as defined above). P<sup>1</sup> can be removed by known techniques. For example, when P<sup>1</sup> is t-butoxycarbonyl group, it can be readily removed by using trifluoroacetic acid or a solution of hydrogen chloride/dioxane. When P<sup>1</sup> is benzyloxycarbonyl group, it can be readily removed by using palladium carbon in combination with hydrogen or ammonium formate. When P<sup>1</sup> is trifluoroacetyl group, it can be readily removed by using an ammonia/methanol solution.

25 [0058]

Process C

[0059]



5 [0060]

Step 1 (Process C)

In this step, the bicycloamide derivative of the general formula (13) (where n and X are as defined above.) and an amine derivative of the formula  $R^1R^2NH$  (where  $R^1$  and  $R^2$  are as defined above) are reacted in the presence of a condensation agent for amidation to give a bicycloamide derivative of the general formula (1) as set forth in claim 1 (where  $R^1$ ,  $R^2$ ,  $P^1$ , n and X are as defined above.). Examples of the condensation agent used in this step include dicyclohexylcarbodiimide (DCC), 10 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), dimethylimidazolinium chloride (DMC), ethyl chloroformate, isobutyl chloroformate and pivaloyl chloride. These agents may be added in the form of solid, liquid or a solution in a proper solvent. When it is desired to use a base 20 in the condensation reaction, the base may be an alkali carbonate, such as sodium bicarbonate and potassium carbonate,

or a tertiary amine, such as triethylamine, diisopropyl ethylamine, N-methylmorpholine, diazabicyclo[5.4.0]-7-undecene, pyridine, 4-dimethylaminopyridine and 1,8-

bis(dimethylamino)naphthalene. The solvent for use in the

condensation reaction may be an inert solvent that does not affect the reaction, such as N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulfoxide, acetonitrile,

tetrahydrofuran, dioxane, ethyl ether, dimethoxyethane, ethyl acetate, toluene and dichloromethane. This condensation

reaction proceeds smoothly at -20 to 150°C. Alternatively, the condensation reaction may be carried out via an active ester or acid chloride having 1-imidazolyl group, 4-nitrophenoxy group, pentafluorophenoxy group, imidoyloxy succinate group or 1-benzotriazolyl group (or 1-benzotriazolyl 3-oxide group).

In such a case, the active ester or acid chloride may be used in the subsequent step after purification or as the unpurified crude product.

[0061]

The advantageous features of the present invention will now be described with reference to experiments and examples, which are not intended to limit the scope of the invention in any way.

[0062]

<Reference Example 1>

Synthesis of 2-tetrahydropyranyl 4-aminobicyclo[2.2.2]octane-

1-carboxylate

[0063]

Step 1:

Synthesis of methyl 4-

5 benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylate

Methyl hydrogen bicyclo[2.2.2]octane-1,4-dicarboxylate (25.0 g), diphenylphosphoryl azide (32.5 g), triethylamine (17.3 mL) and toluene (500 mL) were mixed together. The mixture was stirred for 2 hours at room temperature and was  
10 refluxed for 2 hours. To the resulting mixture, benzylalcohol (122 mL) was added and the mixture was further refluxed for 17 hours. Subsequently, the mixture was allowed to cool and was sequentially washed with a 10% aqueous citric acid, saturated aqueous solution of sodium bicarbonate and saturated brine.  
15 The mixture was then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (eluant: hexane: ethyl acetate = 2:1) to give methyl 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylate (32.2  
20 g).

MS (FAB<sup>+</sup>) m/z: 318 (MH<sup>+</sup>).

[0064]

Step 2:

Synthesis of 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-

25 carboxylic acid

Methyl 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylate (64.3 g) was dissolved in ethanol (1100 mL). To this solution, a 1mol/L aqueous solution of sodium hydroxide (1000 mL) was added and the mixture was stirred at 50°C for 1 hour. Ethanol in the mixture was evaporated under reduced pressure and the residue was washed with diethylether (500 mL), followed by addition of concentrated hydrochloric acid to adjust the pH to 1. The resulting crystals were filtrated, washed with water, dried under reduced pressure to give 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylic acid (56.1 g).

MS (FAB<sup>+</sup>) m/z: 304 (MH<sup>+</sup>).

[0065]

Step 3:

Syntheis of ethyl 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylate

4-Benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylic acid (1.00 g) was dissolved in dichloromethane (10 mL). To this solution, 3,4-dihydro-2H-pyran (1.20 mL) and p-toluenesulfonic monohydrate (6.3 mg) were sequentially added and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was sequentially washed with a saturated aqueous solution of sodium bicarbonate and saturated brine. The mixture was then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified

by silica gel column chromatography (eluant: hexane: ethyl acetate = 4:1) to give 2-tetrahydropyranyl 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylate (1.18 g).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.53-1.95 (m, 18H), 3.67-3.71 (m, 1H), 3.82-3.89 (m, 1H), 4.59 (br, 1H), 5.03 (s, 2H), 5.95 (br, 1H), 7.29-7.38 (m, 5H).

[0066]

Step 4:

10 Synthesis of 2-tetrahydropyranyl 4-aminobicyclo[2.2.2]octane-1-carboxylate

2-Tetrahydropyranyl 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylate (548 mg) was dissolved in ethanol (6 mL). To this solution, 10% palladium-carbon (60 mg) was added and the mixture was stirred at room temperature for 2 hours in a stream of hydrogen. The catalyst in the reaction mixture was filtered through a Celite pad and the filtered catalyst, together with the Celite pad, was washed with ethanol. The filtrate and the washings were combined and concentrated under reduced pressure. The resulting residue was dried under reduced pressure to give 2-tetrahydropyranyl 4-aminobicyclo[2.2.2]octane-1-carboxylate (357 mg).

MS (EI<sup>+</sup>) m/z: 253 (M<sup>+</sup>).

25 [0067]

<Reference Example 2>

Synthesis of 4-aminobicyclo[2.2.2]octane-1-carboxamide

[0068]

Step 1:

5 Synthesis of 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxamide

4-Benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxylic acid (998 mg) was suspended in acetonitrile (20 mL). While the suspension was chilled in an ice bath, N-hydroxybenzotriazole  
10 (605 mg) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (757 mg) were sequentially added. The mixture was stirred at room temperature for 4 hours and was left overnight. Subsequently, 25% aqueous ammonia (1.80 mL) was added while the reaction vessel was chilled in an ice bath.  
15 The mixture was then stirred at room temperature for 1 hour. The insoluble material was filtered and was washed sequentially with acetonitrile and dichloromethane. The filtrate and the washings were combined and concentrated under reduced pressure. The resulting residue was purified by silica  
20 gel chromatography (eluant: ethyl acetate) to give 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxamide (889 mg).

MS (EI<sup>+</sup>) m/z: 302 (M<sup>+</sup>).

[0069]

25 Step 2:

Synthesis of 4-aminobicyclo[2.2.2]octane-1-carboxamide

Using 4-benzyloxycarbonylaminobicyclo[2.2.2]octane-1-carboxamide (367 mg), the same procedure was followed as in Step 4 of Reference Example 1 to give 4-

5 aminobicyclo[2.2.2]octane-1-carboxamide (198 mg).

MS (EI<sup>+</sup>) m/z: 168 (M<sup>+</sup>).

[0070]

<Reference Example 3>

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-(4-

10 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile

[0071]

Step 1:

Synthesis of (2S,4S)-1-[[N-[4-(2-

15 tetrahydropyranyl)oxycarbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

2-Tetrahydropyranyl 4-aminobicyclo[2.2.2]octane-1-carboxylate (62.9 mg) was suspended in acetonitrile (1 mL). To this solution, diisopropylethylamine (47 µL) was added and  
20 (2S,4S)-1-(2-bromoacetyl)-4-fluoropyrrolidine-2-carbonitrile (53.1 mg) in acetonitrile (0.8 mL) was added while the mixture was chilled in an ice bath. The mixture was stirred for 4 hours and concentrated. To the resulting residue, ethyl acetate and water were added, followed by an aqueous sodium  
25 bicarbonate solution to make the pH basic. The solution was



extracted with ethyl acetate. The ethyl acetate layer was washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel chromatography (eluant:

5 dichloromethane: methanol = 10:1) to give (2S,4S)-1-[[N-[4-(2-tetrahydropyranyl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (73.3 mg).

MS (FAB<sup>+</sup>) m/z: 408 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>31</sub>FN<sub>3</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 408.2299; found,

10 408.2295.

[0072]

Step 2:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

15 fluoropyrrolidine-2-carbonitrile

Ethyldiisopropylamine (194 µL) and benzylchloroformate (137 µL) were added dropwise to (2S,4S)-1-[[N-[4-(2-tetrahydropyranyl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (379 mg)

20 in dioxane (5 mL) while the solution was cooled in water. The mixture was stirred at room temperature for 1 hour, followed by addition of 1N hydrochloric acid (0.1 mL) and stirring at room temperature for additional 1 hour. The solvent was evaporated under reduced pressure. The resulting crystal was

25 then washed with diisopropylether and water and was dried

under reduced pressure. The dried crystal was purified by silica gel chromatography (eluant: dichloromethane: methanol = 10:1) to give (2S,4S)-1-[[N-benzyloxycarbonyl-N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

5 fluoropyrrolidine-2-carbonitrile (335 mg).

MS (FAB<sup>+</sup>) m/z: 458 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>29</sub>FN<sub>3</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 458.2091; found, 458.2106.

[0073]

10 <Reference Example 4>

Synthesis of (2S,4S)-1-(2-chloroacetyl)-4-fluoropyrrolidine-2-carbonitrile


According to the process for producing (2S,4S)-1-(2-bromoacetyl)-4-fluoropyrrolidine-2-carbonitrile described in  
15 the publication of WO 02/38541, (2S,4S)-4-fluoropyrrolidine-2-carboxamide hydrochloride (5.00 g) and chloroacetylchloride (2.60 mL) were used to give (2S,4S)-1-(2-chloroacetyl)-4-fluoropyrrolidine-2-carbonitrile (4.96 g).

MS (EI<sup>+</sup>) m/z: 190 (M<sup>+</sup>).

20 HRMS (EI<sup>+</sup>) for C<sub>7</sub>H<sub>8</sub>ClFN<sub>2</sub>O (M<sup>+</sup>): calcd, 190.0309; found, 190.0283.

[Example 1]

[0074]



[0075]

Synthesis of (2S,4S)-1-[[ (4-carbamoylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


4-Aminobicyclo[2.2.2]octane-1-carboxamide (50.0 mg) was  
5 dissolved in N,N-dimethylformamide (2 mL). To this solution,  
potassium carbonate (50.0 mg) was added, followed by dropwise  
addition of (2S,4S)-1-(2-bromoacetyl)-4-fluoropyrrolidine-2-  
carbonitrile (70.0 mg) in N,N-dimethylformamide (1 mL) at room  
temperature. The mixture was stirred at room temperature for  
10 2.5 hours and was concentrated under reduced pressure. The  
resulting residue was purified by silica gel chromatography  
(eluant: chloroform: methanol = 10:1) to give (2S,4S)-1-[[ (4-  
carbamoylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (94.1 mg).

15 MS (FAB<sup>+</sup>) m/z: 323 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>16</sub>H<sub>24</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 323.1883; found,  
323.1903.

[Example 2]

[0076]



20

[0077]

Synthesis of (2S)-1-[[ (4-carbamoylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile


Using 4-aminobicyclo[2.2.2]octane-1-carboxamide (50.0 mg) and (2S)-1-(2-bromoacetyl)pyrrolidine-2-carbonitrile (56.9 mg), the same procedure was followed as in Example 1 to give (2S)-1-[[[4-carbamoylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile (47.5 mg).

MS (FAB<sup>+</sup>) m/z: 305 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>16</sub>H<sub>25</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 305.1798; found, 305.1999.

[Example 3]

10 [0078]



[0079]

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-(4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-Benzyloxycarbonyl-N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (91.5 mg) and 1-hydroxybenzotriazole (45.9 mg) were dissolved in N,N-dimethylformamide (2.0 mL). While the solution was chilled in an ice bath, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (95.9 mg) was added and the mixture was allowed to warm to room temperature and was stirred for 15 hours. The

solvent was evaporated under reduced pressure and the residue was purified by silica gel chromatography (eluant: ethyl acetate) to give (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-

(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-

5 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (92.0 mg).


$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.24-2.25 (m, 12H), 2.57 (t,  $J=15.3$  Hz, 1H), 3.33-4.41 (m, 5H), 4.29-5.50 (m, 4H), 7.30-7.44 (m, 7H), 7.53 (t,  $J=8.0$  Hz, 1H), 8.06 (d,  $J=8.6$  Hz, 1H).

MS ( $\text{FAB}^+$ )  $m/z$ : 575 ( $\text{MH}^+$ ).

10 HRMS ( $\text{FAB}^+$ ) for  $\text{C}_{30}\text{H}_{32}\text{FN}_6\text{O}_5$  ( $\text{MH}^+$ ): calcd, 575.2418; found, 575.2407.

[Example 4]

[0080]



15 [0081]

Synthesis of (2S,4S)-1-[[N-[4-(pyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0082]

20 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(pyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-Benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (20.0 mg) and pyrrolidine (4.4  $\mu$ L) were dissolved in tetrahydrofuran (0.4 mL) and the mixture was stirred at room temperature for 25 minutes. The solvent was removed under reduced pressure and the residue was dissolved in dichloromethane. The organic layer was washed sequentially with 0.1 N aqueous hydrochloric acid, saturated aqueous sodium bicarbonate solution and saturated brine. The organic layer was then dried over anhydrous sodium sulfate and was concentrated under reduced pressure. The resulting residue was purified by silica gel chromatography (eluant: ethyl acetate: methanol = 20:1) to give (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(piperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (16.0 mg).

MS (FAB<sup>+</sup>) m/z: 511 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 511.2721; found, 511.2719.

[0083]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(pyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-Benzyloxycarbonyl-N-[4-(piperidin-1-


yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.0 mg) and 10% palladium carbon (3.0 mg) were dissolved in ethanol (1.0 mL) and dichloromethane (0.5 mL). The mixture was stirred at room temperature for 8 hours in a hydrogen atmosphere. The mixture was then filtered through a Celite pad and the solvent was concentrated under reduced pressure. The resulting residue was purified by column chromatography (eluant: dichloromethane: methanol = 10:1) to give (2S,4S)-1-[[N-[4-(pyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (15.2 mg).

MS (EI<sup>+</sup>) m/z: 376 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>20</sub>H<sub>29</sub>FN<sub>4</sub>O<sub>2</sub> (M<sup>+</sup>): calcd, 376.2275; found, 376.2285.

[Example 5]

[0084]



[0085]

Synthesis of (2S,4S)-1-[[N-[4-(piperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0086]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(piperidin-

1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (100 mg) and piperidine (22.7  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(piperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (81.0 mg).

MS (FAB<sup>+</sup>) m/z: 525 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 525.2877; found, 525.2896.

[0087]

Step 2:

15 Synthesis of (2S,4S)-1-[[N-[4-(piperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-Benzyloxycarbonyl-N-[4-(piperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (20.0 mg) and 10% palladium carbon (12.0 mg) were dissolved in dimethylformamide (0.5 mL). While the solution was chilled in an ice bath, ammonium formate (43.1 mg) was added and the mixture was stirred for 40 minutes at the same temperature. Subsequently, the reaction mixture was filtered through a Celite pad and diluted with



ethyl acetate. The organic layer was washed sequentially with water and saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by column chromatography (eluant:


5 dichloromethane: methanol= 10:1) to give (2S,4S)-1-[[N-[4-(piperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (13.2 mg).

MS (EI<sup>+</sup>) m/z: 390 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>21</sub>H<sub>31</sub>FN<sub>4</sub>O<sub>2</sub> (M<sup>+</sup>): calcd, 390.2431; found, 390.2446.

10 [Example 6]

[0088]



[0089]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(morpholin-4-

15 yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0090]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(morpholin-

20 4-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-

yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and morpholine (9.9  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(morpholin-4-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (43.6 mg).

MS (FAB<sup>+</sup>) m/z: 527 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 527.2670; found, 527.2651.

[0091]

10 Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(morpholin-4-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(morpholin-4-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (34.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(morpholin-4-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (13.2 mg).

MS (FAB<sup>+</sup>) m/z: 393 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 393.2302; found, 393.2304.

[Example 7]

25 [0092]



[0093]

Synthesis of (2S,4S)-1-[[N-[4-(4-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

5 fluoropyrrolidine-2-carbonitrile

[0094]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-

10 ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-

benzyloxycarbonyl-N-[4-(benzotriazol-1-

yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

15 ethoxycarbonylpiperidine (20.1  $\mu$ L) were used to obtain

(2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-

ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (43.1 mg).

MS (FAB<sup>+</sup>) m/z: 597 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>32</sub>H<sub>42</sub>FN<sub>4</sub>O<sub>6</sub> (MH<sup>+</sup>): calcd, 597.3088; found,

597.3096.

[0095]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(4-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.4 mg) was used to obtain (2S,4S)-1-[[N-[4-(4-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (22.1 mg).

MS (FAB<sup>+</sup>) m/z: 463 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 463.2721; found, 463.2723.

[Example 8]

15 [0096]



[0097]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(4-methylpiperazin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

20 [0098]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-

methylpiperazin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-methylpiperazine (14.5  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-methylpiperazin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.0 mg).

MS (FAB<sup>+</sup>) m/z: 540 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>39</sub>FN<sub>5</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 540.2986; found, 540.2974.

[0099]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(4-methylpiperazin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-methylpiperazin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(4-methylpiperazin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.7 mg).

MS (EI<sup>+</sup>) m/z: 405 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>21</sub>H<sub>32</sub>FN<sub>5</sub>O<sub>2</sub> (M<sup>+</sup>): calcd, 405.2540; found, 405.2562.

[Example 9]

[0100]



[0101]

Synthesis of (2S,4S)-1-[[N-[4-(dimethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0102]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(dimethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazole-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2.0 M tetrahydrofuran solution of dimethylamine (65.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(dimethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.2 mg).

MS (FAB<sup>+</sup>) m/z: 485 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 485.2564; found, 485.2554.

[0103]

Step 2:

5 Synthesis of (2S,4S)-1-[[N-[4-(dimethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-

10 (dimethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.0 mg) was used to obtain (2S,4S)-1-[[N-[4-(dimethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (16.7 mg). MS (EI<sup>+</sup>) m/z: 350 (M<sup>+</sup>).

15 HRMS (EI<sup>+</sup>) for C<sub>18</sub>H<sub>27</sub>FN<sub>4</sub>O<sub>2</sub> (M<sup>+</sup>): calcd, 350.2118; found, 350.2156.

[Example 10]

[0104]



[0105]

20 Synthesis of (2S,4S)-1-[[N-[4-[(5-cyanopyridin-2-yl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0106]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[(5-cyanopyridin-2-yl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (5-cyanopyridin-2-yl)piperazine (24.6 mg) were used to obtain (2S,4S)-1-[[benzyloxycarbonyl-N-[4-[(5-cyanopyridin-2-yl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (45.0 mg). MS (FAB<sup>+</sup>) m/z: 628 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>34</sub>H<sub>39</sub>FN<sub>7</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 628.3048; found, 628.3035.

[0107]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-[(5-cyanopyridin-2-yl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[(5-cyanopyridin-2-yl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (45.0 mg) was used to obtain




(2S,4S)-1-[[N-[4-[(5-cyanopyridin-2-yl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (26.1 mg).

MS (FAB<sup>+</sup>) m/z: 494 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>33</sub>FN<sub>7</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 494.2680; found, 494.2684.

[Example 11]

[0108]



10 [0109]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[(2-methoxyphenyl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0110]

15 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[(2-methoxyphenyl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (2-

methoxyphenyl)piperazine (22.9  $\mu$ L) were used to obtain  
(2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[(2-methoxyphenyl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (43.2 mg).

5 MS (FAB<sup>+</sup>) m/z: 632 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>35</sub>H<sub>43</sub>FN<sub>5</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 632.3248; found, 632.3273.

[0111]

Step 2:

10 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[(2-methoxyphenyl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitril


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[(2-methoxyphenyl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (43.0 mg) was used to obtain  
15 (2S,4S)-4-fluoro-1-[[N-[4-[(2-methoxyphenyl)piperazin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (24.0 mg).

20 MS (FAB<sup>+</sup>) m/z: 498 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>37</sub>FN<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 498.2880; found, 498.2905.

[Example 12]

[0112]



[0113]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(4-hydroxypiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0114]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-hydroxypiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-hydroxypiperidine (11.7 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-hydroxypiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (39.0 mg).

MS (FAB<sup>+</sup>) m/z: 541 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 541.2826; found, 541.2836.

[0115]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(4-hydroxypiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-hydroxypiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (39.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(4-hydroxypiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (7.0 mg).

MS (EI<sup>+</sup>) m/z: 406 (M<sup>+</sup>).

HRMS (EI<sup>+</sup>) for C<sub>21</sub>H<sub>31</sub>FN<sub>4</sub>O<sub>3</sub> (M<sup>+</sup>): calcd, 406.2380; found, 406.2399.

[Example 13]

[0116]



[0117]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(4-fluoropiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0118]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-fluoropiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

4-Fluoropiperidine hydrochloride (18.2 mg) was suspended in tetrahydrofuran (0.87 mL). While this suspension was chilled in an ice bath, triethylamine (18.2  $\mu$ L) was added and the mixture was stirred at the same temperature for 35 minutes. Subsequently, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) was added and the mixture was stirred at room temperature overnight. The reaction mixture was diluted with dichloromethane and was washed sequentially with 0.1 N aqueous hydrochloric acid, saturated aqueous sodium bicarbonate solution and saturated brine. The organic layer was dried over anhydrous sodium sulfate and was concentrated under reduced pressure. The resulting residue was purified by column chromatography (eluant: ethyl acetate: methanol = 20:1) to give (2S,4S)-1-[[N-benzyloxycarbonyl-[4-(4-fluoropiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (37.0 mg).

MS (FAB<sup>+</sup>) m/z: 543 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>37</sub>F<sub>2</sub>N<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 543.2783; found, 543.2794.

[0119]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(4-fluoropiperidin-1-

yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-fluoropiperidin-1-


5 yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (37.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(4-fluoropiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (16.4 mg).

10 MS (FAB<sup>+</sup>) m/z: 409 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>31</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 409.2415; found, 409.2392.

[Example 14]

[0120]



15

[0121]

Synthesis of (2S,4S)-1-[[N-[4-(4-benzylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

20 [0122]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-benzylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-benzylpiperidine (22.9  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-benzylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (45.9 mg).

10 MS (FAB<sup>+</sup>) m/z: 615 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>36</sub>H<sub>44</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 615.3347; found, 615.3388.

[0123]

Step 2:

15 Synthesis of (2S,4S)-1-[[N-[4-(4-benzylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(4-benzylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (45.9 mg) was used to obtain (2S,4S)-1-[[N-[4-(4-benzylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (23.0 mg).

25 MS (FAB<sup>+</sup>) m/z: 481 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 481.2979; found, 481.2935.

[Example 15]

[0124]



[0125]

Synthesis of (2S,4S)-1-[[[4-(azetidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0126]

Step 1:

Synthesis of (2S,4S)-1-[[N-[4-(azetidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and azetidine hydrochloride (12.2 mg) were used to obtain (2S,4S)-1-[[N-[4-(azetidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (32.0 mg).



MS (FAB<sup>+</sup>) m/z: 497 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 497.2564; found, 497.2567.

[0127]

5 Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(azetidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


10 In a similar manner to Example 5, (2S,4S)-1-[[N-[4-(azetidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (27.0 mg) was used to obtain (2S,4S)-1-[[N-[4-(azetidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (10.0 mg).

15 MS (FAB<sup>+</sup>) m/z: 363 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 363.2196; found, 363.2221.

[Example 16]

[0128]



20

[0129]

Synthesis of (2S,4S,3'R)-4-fluoro-1-[[N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0130]

Step 1:

Synthesis of (2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-

5 fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
10 fluoropyrrolidine-2-carbonitrile (50.0 mg) and (3R)-3-fluoropyrrolidine hydrochloride (16.4 mg) were used to obtain (2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (39.7 mg).

15 MS (FAB<sup>+</sup>) m/z: 529 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>35</sub>F<sub>2</sub>N<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 529.2626; found, 529.2642.

[0131]

Step 2:

20 Synthesis of (2S,4S,3'R)-4-fluoro-1-[[N-[4-(3-  
fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

25


fluoropyrrolidine-2-carbonitrile (39.7 mg) was used to obtain (2S,4S,3'R)-4-fluoro-1-[[N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.8 mg).

5 MS (FAB<sup>+</sup>) m/z: 395 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>29</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 395.2259; found, 395.2216.

[Example 17]

[0132]



10

[0133]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(3,3-difluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

15 [0134]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(3,3-difluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

20 In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 3,3-

difluoropyrrolidine hydrochloride (18.7 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(3,3-difluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (40.8 mg).

5 MS (FAB<sup>+</sup>) m/z: 547 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>34</sub>F<sub>3</sub>N<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 547.2532; found, 547.2549.

[0135]

Step 2:

10 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(3,3-difluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(3,3-difluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (40.8 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(3,3-difluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (26.6 mg).

20 MS (FAB<sup>+</sup>) m/z: 413 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>28</sub>F<sub>3</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 413.2164; found, 413.2126.

[Example 18]

[0136]



[0137]

Synthesis of (2S,4S,3'S)-4-fluoro-1-[[N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0138]

Step 1:

Synthesis of (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (3S)-3-fluoropyrrolidine hydrochloride (16.4 mg) were used to obtain (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (35.3 mg).

MS (FAB<sup>+</sup>) m/z: 529 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>35</sub>F<sub>2</sub>N<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 529.2626; found, 529.2642.

[0139]

Step 2:

Synthesis of (2S,4S,3'S)-4-fluoro-1-[[N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (35.3 mg) was used to obtain (2S,4S,3'S)-4-fluoro-1-[[N-[4-(3-fluoropyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (19.9 mg).

MS (FAB<sup>+</sup>) m/z: 395 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>29</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 395.2259; found, 395.2266.

[Example 19]

[0140]



[0141]

Synthesis of (2S,4S,3'S)-1-[[N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0142]

Step 1:

(2S,4S,3'S)-1-[[N-Benzyloxycarbonyl-N-[4-(3-

ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (70.0 mg) and (S)-(+)-nipecotic acid ethyl ester (28.0  $\mu$ L) were used to obtain (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (53.8 mg). MS (FAB<sup>+</sup>) m/z: 597 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>32</sub>H<sub>42</sub>FN<sub>4</sub>O<sub>6</sub> (MH<sup>+</sup>): calcd, 597.3088; found, 597.3108.

[0143]

Step 2:

Synthesis of (2S,4S,3'S)-1-[[N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (53.8 mg) was used to obtain (2S,4S,3'S)-1-[[N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (25.2 mg).

MS (FAB<sup>+</sup>) m/z: 463 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 463.2721; found, 463.2690.

[Example 20]

5 [0144]



[0145]

Synthesis of (2S,4S,3'R)-1-[[N-[4-(3-ethoxycarbonylpiperidin-  
1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

10 fluoropyrrolidine-2-carbonitrile

[0146]

Step 1:

Synthesis of (2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-  
ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

15 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-

yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (80.0 mg) and (R)-(-)-

20 nipecotic acid ethyl ester (32.2  $\mu$ L) were used to obtain

(2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-

ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (78.3 mg).



MS (FAB<sup>+</sup>) m/z: 597 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>32</sub>H<sub>42</sub>FN<sub>4</sub>O<sub>6</sub> (MH<sup>+</sup>): calcd, 597.3088; found, 597.3096.

[0147]

5 Step 2:

Synthesis of (2S,4S,3'R)-1-[[N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Exemplar 5, (2S,4S,3'R)-1-[[N-  
10 benzyloxycarbonyl-N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (78.3 mg) was used to obtain (2S,4S,3'R)-1-[[N-[4-(3-ethoxycarbonylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
15 fluoropyrrolidine-2-carbonitrile (47.2 mg).

MS (FAB<sup>+</sup>) m/z: 463 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 463.2721; found, 463.2711.

[Example 21]

20 [0148]



[0149]

Synthesis of (2S,4S,3'S)-1-[[N-[4-(3-hydroxypyrrolidin-1-

yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile

[0150]

Step 1:

5 Synthesis of (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-  
hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (3S)-3-hydroxypyrrolidine (9.1  $\mu$ L) were used to obtain (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (35.0 mg).

MS (FAB<sup>+</sup>) m/z: 527 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 527.2670; found, 527.2679

[0151]

20 Step 2:

Synthesis of (2S,4S,3'S)-1-[[N-[4-(3-hydroxypyrrolidin-1-  
yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-(3-hydroxypyrrolidin-1-


yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (35.0 mg) was used to obtain (2S,4S,3'S)-1-[[N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (15.2 mg).

MS (FAB<sup>+</sup>) m/z: 393 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 393.2302; found, 393.2300.

[Example 22]

10 [0152]



[0153]

Synthesis of (2S,4S,3'R)-1-[[N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

15 fluoropyrrolidine-2-carbonitrile

[0154]

Step 1:

Synthesis of (2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-

20 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (80.0 mg) and (3R)-3-hydroxypyrrolidine (16.9  $\mu$ L) were used to obtain (2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

5 fluoropyrrolidine-2-carbonitrile (75.0 mg).

MS (FAB<sup>+</sup>) m/z: 527 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 527.2670; found, 527.2679

[0155]

10 Step 2:

Synthesis of (2S,4S,3'R)-1-[[N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S,3'R)-1-[[N-benzyloxycarbonyl-N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (75.0 mg) was used to obtain (2S,4S,3'R)-1-[[N-[4-(3-hydroxypyrrolidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (38.9 mg).

MS (FAB<sup>+</sup>) m/z: 393 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 393.2302; found, 393.2274.

[Example 23]

25 [0156]



[0157]

Synthesis of (2S,4S)-1-[[[4-(4-acetylamino-4-phenylpiperidin-  
1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
 5 fluoropyrrolidine-2-carbonitrile

[0158]

Step 1:

Synthesis of (2S,4S)-1-[[N-[4-(4-acetylamino-4-  
phenylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl-N-  
 10 benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-  
carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-  
 benzyloxycarbonyl-N-[4-(benzotriazol-1-  
 yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
 15 fluoropyrrolidine-2-carbonitrile (80.0 mg) and 4-acetylamino-  
 4-phenylpiperidine hydrochloride (53.2 mg) were used to obtain  
 (2S,4S)-1-[[N-[4-(4-acetylamino-4-phenylpiperidin-1-  
 yl)carbonylbicyclo[2.2.2]oct-1-yl-N-  
 benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-  
 20 carbonitrile (64.5 mg).

MS (FAB<sup>+</sup>) m/z: 658 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>37</sub>H<sub>45</sub>FN<sub>5</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 658.3405; found,

658.3414.

[0159]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(4-acetylamino-4-

5 phenylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-[4-(4-acetylamino-4-phenylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-  
10 carbonitrile (64.5 mg) was used to obtain (2S,4S)-1-[[N-[4-(4-acetylamino-4-phenylpiperidin-1-yl)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.2 mg).

MS (FAB<sup>+</sup>) m/z: 524 (MH<sup>+</sup>).

15 HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>39</sub>FN<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 524.3037; found, 524.3047.

[Example 24]

[0160]



20 [0161]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
methylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0162]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-  
methylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
5 fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and methylamine  
10 (2.0 mol/l THF solution, 60.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-methylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (28.9 mg).  
MS (FAB<sup>+</sup>) m/z: 471 (MH<sup>+</sup>).  
15 Rf 0.25 (ethyl acetate: methanol = 9:1).

[0163]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
methylamino)carbonylbicyclo[2.2.2]oct-1-  
20 yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-methylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (27.0 mg) was used to obtain  
25 (2S,4S)-4-fluoro-1-[[N-[4-(N-

methyamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (10.8 mg).


MS (FAB<sup>+</sup>) m/z: 337 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>17</sub>H<sub>25</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 337.2040; found,

5 337.2040.

[Example 25]

[0164]



[0165]

10 Synthesis of (2S,4S)-1-[[N-[4-(N-ethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0166]

Step 1:

15 Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-ethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethylamine (2.0 mol/L THF solution, 60.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-



ethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (24.6 mg).

MS (FAB<sup>+</sup>) m/z: 485 (MH<sup>+</sup>).

Rf 0.33 (ethyl acetate: methanol = 15:1).

5 [0167]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(N-

ethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile

10 In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-ethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (22.6 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-


15 ethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (11.4 mg).

MS (FAB<sup>+</sup>) m/z: 351 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>18</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 351.2196; found, 351.2181.

20 [Example 26]

[0168]



[0169]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-propylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile  
[0170]

5 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-propylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and propylamine (10.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-propylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (28.7 mg).  
MS (FAB<sup>+</sup>) m/z: 499 (MH<sup>+</sup>).  
Rf 0.38 (ethyl acetate: methanol = 15:1).  
[0171]

Step 2:

20 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-propylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-propylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-


fluoropyrrolidine-2-carbonitrile (25.8 mg) was used to obtain  
(2S,4S)-4-fluoro-1-[[N-[4-(N-  
propylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.9 mg).

5 MS (FAB<sup>+</sup>) m/z: 365 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 365.2353; found,  
365.2382.

[Example 27]

[0172]



10

[0173]

Synthesis of (2S,4S)-1-[[N-[4-(N-  
cyclopropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-  
4-fluoropyrrolidine-2-carbonitrile

15 [0174]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-  
cyclopropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-  
4-fluoropyrrolidine-2-carbonitrile

20 In a similar manner to Example 4, (2S,4S)-1-[[N-  
benzyloxycarbonyl-N-[4-(benzotriazol-1-  
yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (50.0 mg) and

cyclopropylamine (8.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-cyclopropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.6 mg).

5 MS (FAB<sup>+</sup>) m/z: 497 (MH<sup>+</sup>).

Rf 0.35 (ethyl acetate: methanol = 15:1).

[0175]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(N-cyclopropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-cyclopropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.1 mg) was used to obtain (2S,4S)-1-[[N-[4-(N-cyclopropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (16.7 mg).

MS (FAB<sup>+</sup>) m/z: 363 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 363.2196; found, 363.2217.

[Example 28]

[0176]



[0177]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-1-methylethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0178]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 1-methylethylamine (10.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (33.6 mg).

MS (FAB<sup>+</sup>) m/z: 499 (MH<sup>+</sup>).

Rf 0.25 (ethyl acetate: methanol = 20:1).

[0179]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-1-

methylethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-


5 methylethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.2 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-1-methylethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.7 mg).

10 MS (FAB<sup>+</sup>) m/z: 365 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 365.2353; found, 365.2345.

[Example 29]

[0180]



15

[0181]

Synthesis of (2S,4S,1'RS)-4-fluoro-1-[[N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

20 [0182]

Step 1:

Synthesis of (2S,4S,1'RS)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 1-methylpropylamine (12.0  $\mu$ L) were used to obtain (2S,4S,1'RS)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (32.0 mg).

10 MS (FAB<sup>+</sup>) m/z: 513 (MH<sup>+</sup>).

Rf 0.33 (ethyl acetate).

[0183]

Step 2:

Synthesis of (2S,4S,1'RS)-4-fluoro-1-[[N-[4-(N-1-

15 methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S,1'RS)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (29.0 mg) was used to obtain (2S,4S,1'RS)-4-fluoro-1-[[N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.9 mg).


MS (FAB<sup>+</sup>) m/z: 379 (MH<sup>+</sup>).

25 HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 379.2509; found,

379.2497.

[Example 30]

[0184]



5 [0185]

Synthesis of (2S,4S)-1-[[N-[4-(N-2,2-  
dimethylethylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0186]

10 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-2,2-  
dimethylethylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-  
15 benzyloxycarbonyl-N-[4-(benzotriazol-1-  
yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2,2-  
dimethylethylamine (12.0  $\mu$ L) were used to obtain (2S,4S)-1-  
[[N-benzyloxycarbonyl-N-[4-(N-2,2-  
20 dimethylethylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.5 mg).  
MS (FAB<sup>+</sup>) m/z: 513 (MH<sup>+</sup>).  
Rf 0.45 (ethyl acetate).



[0187]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(N-2,2-

dimethylethylamino)carbonylbicyclo[2.2.2]oct-1-

5 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-2,2-

dimethylethylamino)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (28.9 mg)

10 was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-2,2-

dimethylethylamino)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile (17.3 mg).


MS (FAB<sup>+</sup>) m/z: 379 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 379.2509; found,

15 379.2518.

[Example 31]

[0188]



[0189]

20 Synthesis of (2S,4S)-1-[[N-[4-(N-

cyclohexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-

4-fluoropyrrolidine-2-carbonitrile

[0190]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-cyclohexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

5           In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and cyclohexylamine (13.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-cyclohexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (35.0 mg).  
10 MS (FAB<sup>+</sup>) m/z: 539 (MH<sup>+</sup>).  
Rf 0.35 (ethyl acetate).

[0191]

15 Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(N-cyclohexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


          In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-cyclohexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.6 mg) was used to obtain (2S,4S)-1-[[N-[4-(N-cyclohexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (17.5 mg).  
20  
25

MS (FAB<sup>+</sup>) m/z: 405 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 405.2666; found, 405.2628.

[Example 32]

5 [0192]



[0193]

Synthesis of (2S,4S)-1-[[N-[4-(N-benzylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
10 fluoropyrrolidine-2-carbonitrile

[0194]

Step 1:

Synthesis of (2S,4S)-1-[[N-[4-(N-benzylamino)carbonylbicyclo[2.2.2]oct-1-yl]-N-  
15 benzyloxycarbonylamino]acetyl]-4-fluoropyrrolidine-2-  
carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

20 fluoropyrrolidine-2-carbonitrile (50.0 mg) and benzylamine (13.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-[4-(N-benzylamino)carbonylbicyclo[2.2.2]oct-1-yl]-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-

carbonitrile (32.1 mg).

MS (FAB<sup>+</sup>) m/z: 547 (MH<sup>+</sup>).

Rf 0.30 (ethyl acetate).

[0195]

5 Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(N-benzylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


10 In a similar manner to Example 5, (2S,4S)-1-[[N-[4-(N-benzylamino)carbonylbicyclo[2.2.2]oct-1-yl-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.0 mg) was used to obtain (2S,4S)-1-[[N-[4-(N-benzylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (13.7 mg).

15 MS (FAB<sup>+</sup>) m/z: 413 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 413.2353; found, 413.2345.

[Example 33]

[0196]



20

[0197]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-phenylamino)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0198]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-

5 phenylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and aniline (10.0  
10  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-phenylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (35.4 mg).

MS (FAB<sup>+</sup>) m/z: 533 (MH<sup>+</sup>).

15 R<sub>f</sub> 0.33 (ethyl acetate: hexane = 4:1).

[0199]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-

phenylamino)carbonylbicyclo[2.2.2]oct-1-  
20 yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-phenylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.2 mg) was used to obtain  
25 (2S,4S)-4-fluoro-1-[[N-[4-(N-

phenylamino)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile (16.6 mg).


MS (FAB<sup>+</sup>) m/z: 399 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 399.2196; found,

5 399.2220.

[Example 34]

[0200]



10 [0201]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-3-  
hydroxyadamantylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0202]

15 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-3-  
hydroxyadamantylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-  
20 benzyloxycarbonyl-N-[4-(benzotriazol-1-  
yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (50.0 mg) and 3-  
aminoadamantanol (18.9 mg) were used to obtain (2S,4S)-1-[[N-

benzyloxycarbonyl-N-[4-(N-3-hydroxyadamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (40.0 mg).  
MS (FAB<sup>+</sup>) m/z: 607 (MH<sup>+</sup>).

5 Rf 0.33 (ethyl acetate: methanol = 9:1).

[0203]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-3-hydroxyadamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


10

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-3-hydroxyadamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (37.4 mg)  
15 was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-3-hydroxyadamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (21.9 mg).  
MS (FAB<sup>+</sup>) m/z: 473 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 473.2928; found,  
20 473.2952.

[Example 35]

[0204]



[0205]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-2-hydroxyethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

5 [0206]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-2-hydroxyethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

10 In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-aminoethanol (6.9 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-2-hydroxyethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (27.2 mg).  
15 MS (FAB<sup>+</sup>) m/z: 501 (MH<sup>+</sup>).

Rf 0.31 (dichloromethane: methanol = 15:1).

[0207]

20 Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-2-hydroxyethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-2-

25



hydroxyethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (25.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-2-hydroxyethylamino)carbonylbicyclo[2.2.2]oct-1-


5 yl]amino]acetyl]pyrrolidine-2-carbonitrile (12.2 mg).

MS (FAB<sup>+</sup>) m/z: 367 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>18</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 367.2145; found, 367.2150.

[Example 36]

10 [0208]



[0209]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-3-pyridylmethylamino)carbonylbicyclo[2.2.2]oct-1-

15 yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0210]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-3-pyridylmethylamino)carbonylbicyclo[2.2.2]oct-1-

20 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and 3-pyridylmethylaniline (12.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-3-pyridylmethylaniline)carbonylbicyclo[2.2.2]oct-1-

5 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.2 mg).

MS (FAB<sup>+</sup>) m/z: 548 (MH<sup>+</sup>).

R<sub>f</sub> 0.33 (dichloromethane:methanol=15:1).

[0211]

Step 2:

10 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-3-pyridylmethylaniline)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-3-


15 pyridylmethylaniline)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (34.6 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-3-pyridylmethylaniline)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.0 mg).

20 MS (FAB<sup>+</sup>) m/z: 414 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>29</sub>FN<sub>5</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 414.2305; found, 414.2311.

[Example 37]

[0212]



[0213]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-4-fluorobenzylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0214]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-4-fluorobenzylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-fluorobenzylamine (13.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-4-fluorobenzylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (38.3 mg).

MS (FAB<sup>+</sup>) m/z: 565 (MH<sup>+</sup>).

Rf 0.48 (ethyl acetate: methanol = 20:1).

[0215]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-4-

fluorobenzylamino) carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-4-


5 fluorobenzylamino) carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (35.3 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-4-fluorobenzylamino) carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (17.3 mg).

10 MS (FAB<sup>+</sup>) m/z: 431 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>29</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 431.2259; found, 431.2246.

[Example 38]

[0216]



15

[0217]

Synthesis of (2S,4S)-1-[[N-[4-(N-diphenylmethylamino) carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

20 [0218]

Step 1:

Synthesis of (2S,4S)-1-[[N-[4-(N-

diphenylmethylamino)carbonylbicyclo[2.2.2]oct-1-yl]-N-  
benzyloxycarbonylamino]acetyl]-4-fluoropyrrolidine-2-  
carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-  
5 benzyloxycarbonyl-N-[4-(benzotriazol-1-  
yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (50.0 mg) and  
diphenylmethylamine (20.0  $\mu$ L) were used to obtain (2S,4S)-1-  
[[N-[4-(N-diphenylmethylamino)carbonylbicyclo[2.2.2]oct-1-yl]-  
10 N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-  
carbonitrile (40.0 mg).

MS (FAB<sup>+</sup>) m/z: 623 (MH<sup>+</sup>).

Rf 0.63 (ethyl acetate).

[0219]

15 Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(N-  
diphenylmethylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-[4-(N-  
20 diphenylmethylamino)carbonylbicyclo[2.2.2]oct-1-yl-N-  
benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-  
carbonitrile (37.4 mg) was used to obtain (2S,4S)-1-[[N-[4-(N-  
diphenylmethylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (20.3 mg).

25 MS (FAB<sup>+</sup>) m/z: 489 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 489.2666; found, 489.2675.

[Example 39]

[0220]



[0221]

Synthesis of (2S,4S,3'S)-1-[[N-[4-[N-(1-benzylpyrrolidin-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0222]

Step 1:

Synthesis of (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(1-benzylpyrrolidin-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 3(S)-amino-1-benzylpyrrolidine (20.0  $\mu$ L) were used to obtain (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(1-benzylpyrrolidin-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.9 mg).

MS (FAB<sup>+</sup>) m/z: 616 (MH<sup>+</sup>).

Rf 0.25 (dichloromethane: methanol = 20:1).

[0223]

Step 2:

5 Synthesis of (2S,4S,3'S)-1-[[N-[4-[N-(1-benzylpyrrolidin-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, ((2S,4S,3'S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(1-benzylpyrrolidin-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (34.5 mg) was used to obtain  
10 (2S,4S,3'S)-1-[[N-[4-[N-(1-benzylpyrrolidin-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (11.7 mg).

MS (FAB<sup>+</sup>) m/z: 482 (MH<sup>+</sup>).

15 HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>37</sub>FN<sub>5</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 482.2931; found, 482.2926.

[Example 40]

[0224]



20 [0225]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-2-fluoroethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0226]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-2-  
fluoroethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-  
5 4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-  
10 fluoroethylamine hydrochloride (11.2 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-2-fluoroethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (33.9 mg).  
MS (FAB<sup>+</sup>) m/z: 503 (MH<sup>+</sup>).  
15 Rf 0.33 (ethyl acetate: methanol = 15:1).

[0227]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-2-  
fluoroethylamino)carbonylbicyclo[2.2.2]oct-1-  
20 yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-4-fluorobenzylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (32.0 mg)  
25 was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-2-



fluoroethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.2 mg).


MS (FAB<sup>+</sup>) m/z: 369 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>18</sub>H<sub>27</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 369.2102; found,

5 369.2103.

[Example 41]

[0228]



[0229]

10 Synthesis of (2S,4S)-1-[[N-[4-(N-2-cyanoethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0230]

Step 1:

15 Synthesis of (2S,4S)-1-[[N-[4-(N-2-cyanoethylamino)carbonylbicyclo[2.2.2]oct-1-yl]-N-benzyloxycarbonylamino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-cyanoethylamine (9.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-[4-

(N-2-cyanoethylamino)carbonylbicyclo[2.2.2]oct-1-yl-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (29.1 mg).

MS (FAB<sup>+</sup>) m/z: 510 (MH<sup>+</sup>).

5 Rf 0.40 (ethyl acetate: methanol = 9:1).

[0231]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-(N-2-  
cyanoethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-  
10 4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-[4-(N-2-cyanoethylamino)carbonylbicyclo[2.2.2]oct-1-yl-N-benzyloxycarbonyl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (25.2 mg) was used to obtain (2S,4S)-1-[[N-[4-(N-


15 2-cyanoethylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (14.5 mg).

MS (FAB<sup>+</sup>) m/z: 376 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>27</sub>FN<sub>5</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 376.2149; found, 376.2161.

20 [Example 42]

[0232]



[0233]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-butylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile  
[0234]

5 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-butylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and butylamine (11.5  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-butylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (44.0 mg).  
MS (FAB<sup>+</sup>) m/z: 513 (MH<sup>+</sup>).  
Rf 0.25 (ethyl acetate).  
[0235]

Step 2:

20 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-butylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-butylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-


fluoropyrrolidine-2-carbonitrile (37.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-butylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (22.8 mg).

5 MS (FAB<sup>+</sup>) m/z: 379 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 379.2509; found, 379.2504.

[Example 43]

[0236]



10

[0237]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-pentylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

15 [0238]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-pentylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

20

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and pentylamine

(15.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-pentylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (39.6 mg).  
MS (FAB<sup>+</sup>) m/z: 527 (MH<sup>+</sup>).

5 Rf 0.43 (ethyl acetate: methanol = 20:1).

[0239]

Step 2:


Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
pentylamino)carbonylbicyclo[2.2.2]oct-1-  
10 yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-pentylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (37.6 mg) was used to obtain  
15 (2S,4S)-4-fluoro-1-[[N-[4-(N-pentylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (21.4 mg).  
MS (FAB<sup>+</sup>) m/z: 393 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 393.2666; found,  
20 393.2633.

[Example 44]

[0240]



[0241]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-hexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

5 [0242]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-hexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

10 In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and hexylamine (15.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-hexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (42.7 mg).  
MS (FAB<sup>+</sup>) m/z: 541 (MH<sup>+</sup>).

Rf 0.45 (ethyl acetate:methanol=20:1).

[0243]

20 Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-hexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-

hexylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (41.0 mg) was used to obtain

(2S,4S)-4-fluoro-1-[[N-[4-(N-

hexylamino)carbonylbicyclo[2.2.2]oct-1-


5 yl]amino]acetyl]pyrrolidine-2-carbonitrile (24.5 mg).

MS (FAB<sup>+</sup>) m/z: 407 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 407.2822; found, 407.2794.

[Example 45]

10 [0244]



[0245]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
heptylamino)carbonylbicyclo[2.2.2]oct-1-

15 yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0246]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-  
heptylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

20 fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and heptylamine (20.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-heptylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (42.9 mg).

5 MS (FAB<sup>+</sup>) m/z: 555 (MH<sup>+</sup>).

Rf 0.45 (ethyl acetate).

[0247]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
10 heptylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-heptylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
15 fluoropyrrolidine-2-carbonitrile (39.7 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-heptylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (22.2 mg).


MS (FAB<sup>+</sup>) m/z: 421 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 421.2979; found, 421.3002.

[Example 46]

[0248]





[0249]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
octylamino)carbonylbicyclo[2.2.2]oct-1-

5 yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0250]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-  
octylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

10 fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and octylamine (15.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-octylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (42.4 mg). MS (FAB<sup>+</sup>) m/z: 569 (MH<sup>+</sup>).

R<sub>f</sub> 0.50 (ethyl acetate: methanol = 20:1).

20 [0251]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
octylamino)carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-octylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

5 fluoropyrrolidine-2-carbonitrile (41.1 mg) was used to obtain

(2S,4S)-4-fluoro-1-[[N-[4-(N-

octylamino)carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (17.2 mg).

MS (FAB<sup>+</sup>) m/z: 435 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>40</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 435.3135; found, 435.3160.

[Example 47]

[0252]



15 [0253]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-hydroxybicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0254]

20 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(4-hydroxybicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-aminobicyclo[2.2.2]octane-1-ol (13.5 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(4-hydroxybicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.2 mg).

MS (FAB<sup>+</sup>) m/z: 581 (MH<sup>+</sup>).

Rf 0.38 (ethyl acetate: methanol = 9:1).

[0255]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-hydroxybicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(4-hydroxybicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (28.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-hydroxybicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (11.9 mg).


MS (FAB<sup>+</sup>) m/z: 447 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 447.2771; found,

447.2798.

[Example 48]

[0256]



5 [0257]

Synthesis of (2S,4S,1'S)-4-fluoro-1-[[N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0258]

10 Step 1:

Synthesis of (2S,4S,1'S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (350.0 mg) and 1(S)-methylpropylamine (80.0  $\mu$ L) were used to obtain (2S,4S,1'S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (296.9 mg).

MS (FAB<sup>+</sup>) m/z: 513 (MH<sup>+</sup>).

Rf 0.38 (ethyl acetate: methanol = 20:1).

[0259]

Step 2:

Synthesis of (2S,4S,1'S)-4-fluoro-1-[[N-[4-(N-1-  
methylpropylamino)carbonylbicyclo[2.2.2]oct-1-  
5 yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S,1'S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (294.0 mg)  
10 was used to obtain (2S,4S,1'S)-4-fluoro-1-[[N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (172.4 mg).

MS (FAB<sup>+</sup>) m/z: 379 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 379.2509; found,  
15 379.2469.

[Example 49]

[0260]



[0261]

20 Synthesis of (2S,4S,1'R)-4-fluoro-1-[[N-[4-(N-1-  
methylpropylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0262]

Step 1:

Synthesis of (2S,4S,1'R)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

5           In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (350.0 mg) and 1(R)-methylpropylamine (80.0μL) were used to obtain (2S,4S,1'R)-1-  
10   [[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (292.4 mg).  
MS (FAB<sup>+</sup>) m/z: 513 (MH<sup>+</sup>).  
Rf 0.38 (ethyl acetate: methanol = 20:1).  
15   [0263]

Step 2:

Synthesis of (2S,4S,1'R)-4-fluoro-1-[[N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

20           In a similar manner to Example 5, (2S,4S,1'R)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (290.0 mg) was used to obtain (2S,4S,1'R)-4-fluoro-1-[[N-[4-(N-1-methylpropylamino)carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (158.3 mg).

MS (FAB<sup>+</sup>) m/z: 379 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 379.2509; found, 379.2477.

5 [Example 50]

[0264]



[0265]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(thiazol-2-

10 yl]amino]carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-(4-Carboxybicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.0 mg),

along with 1-hydroxybenzotriazole, was dissolved in N,N-

15 dimethylformamide (1.0 mL). To this solution, 2-aminothiazole

(18.6 mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide

hydrochloride (53.4 mg) were added, and the mixture was

stirred at room temperature for 15 hours. The solvent was

evaporated under reduced pressure and the resulting residue

20 was purified by preparative thin-layer chromatography

(solvent: dichloromethane: methanol = 9:1) to give (2S,4S)-4-

fluoro-1-[[N-[4-[N-(thiazol-2-

yl]amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (17.4 mg).

MS (FAB<sup>+</sup>) m/z: 406 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>25</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 406.1713; found, 406.1695.

5 [Example 51]

[0266]



[0267]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-

10 fluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-(4-Carboxybicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.0 mg),  
along with 1-hydroxybenzotriazole, was dissolved in N,N-

15 dimethylformamide (1.0 mL). While the solution was chilled in  
an ice bath, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide  
hydrochloride (53.4 mg) was added and the mixture was allowed  
to warm to room temperature and was stirred for 1 hour.

Subsequently, 4-fluoroaniline (17.8  $\mu$ L) was added and the

20 mixture was stirred for additional 2 hours. The solvent was  
evaporated under reduced pressure and the resulting residue  
was purified by preparative thin-layer chromatography

(solvent: dichloromethane: methanol = 4:1) to give (2S,4S)-4-




fluoro-1-[[N-[4-[N-(4-fluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.6 mg).

MS (FAB<sup>+</sup>) m/z: 417 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>27</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 417.2102; found, 417.2078.

[Example 52]

[0268]



10 [0269]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-propenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 51, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.0 mg) and allylamine (14.0  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-propenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.7 mg).

20 MS (FAB<sup>+</sup>) m/z: 363 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 363.2196; found, 363.2172.

[Example 53]

[0270]



[0271]

Synthesis of (2S,4S,3'S)-4-fluoro-1-[[N-[4-[N-(2-oxo-1-  
5 azacyclohept-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0272]

Step 1:

Synthesis of (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-[4-[N-(2-oxo-  
10 1-azacyclohept-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (S)-5-amino-ε-caprolactam hydrochloride (18.6 mg) were used to obtain (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-[4-[N-(2-oxo-1-azacyclohept-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.3 mg).  
20 MS (FAB<sup>+</sup>) m/z: 568 (MH<sup>+</sup>).

R<sub>f</sub> 0.38 (ethyl acetate: methanol = 5:1).

[0273]

Step 2:

Synthesis of (2S,4S,3'S)-4-fluoro-1-[[N-[4-[N-(2-oxo-1-azacyclohept-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-[4-[N-(2-oxo-1-azacyclohept-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (28.2 mg) was used to obtain (2S,4S,3'S)-4-fluoro-1-[[N-[4-[N-(2-oxo-1-azacyclohept-3-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (13.0 mg).

MS (FAB<sup>+</sup>) m/z: 434 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>33</sub>FN<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 434.2567; found, 434.2566.

[Example 54]

[0274]



[0275]

Synthesis of (2S,4S,1'S)-4-fluoro-1-[[N-[4-[N-(1-ethoxycarbonyl-2-phenylethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0276]

Step 1:

Synthesis of (2S,4S,1'S)-1-[[N-benzyloxycarbonyl-[4-[N-(1-

ethoxycarbonyl-2-phenylethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and L-phenylalanine ethyl ester hydrochloride (26.0 mg) were used to obtain (2S,4S,1'S)-1-[[N-benzyloxycarbonyl-[4-[N-(1-ethoxycarbonyl-2-phenylethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (55.0 mg). MS (FAB<sup>+</sup>) m/z: 633 (MH<sup>+</sup>). Rf 0.48 (ethyl acetate).

[0277]

Step 2:

15 Synthesis of (2S,4S,1'S)-1-[[N-[4-[N-(1-ethoxycarbonyl-2-phenylethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S,3'S)-1-[[N-benzyloxycarbonyl-[4-[N-(1-ethoxycarbonyl-2-phenylethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (53.0 mg) was used to obtain (2S,4S,3'S)-1-[[N-[4-[N-(1-ethoxycarbonyl-2-phenylethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (16.0 mg).

25 MS (FAB<sup>+</sup>) m/z: 499 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 499.2721; found, 499.2729.

[Example 55]

[0278]



[0279]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-ethoxycarbonylbicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0280]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(4-ethoxycarbonylbicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethyl 4-aminobicyclo[2.2.2]octane-1-carboxylate (22.3 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(4-ethoxycarbonylbicyclo[2.2.2]oct-1-

yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (40.7 mg).

MS (FAB<sup>+</sup>) m/z: 637 (MH<sup>+</sup>).

Rf 0.40 (ethyl acetate).

5 [0281]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-

ethoxycarbonylbicyclo[2.2.2]oct-1-

yl)amino]carbonylbicyclo[2.2.2]oct-1-

10 yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-(4-ethoxycarbonylbicyclo[2.2.2]oct-1-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (38.7 mg) was used to obtain

15 (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-

ethoxycarbonylbicyclo[2.2.2]oct-1-

yl)amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (22.3 mg).

MS (FAB<sup>+</sup>) m/z: 503 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>40</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 503.3034; found, 503.3080.

[Example 56]

[0282]



[0283]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[4-(piperidin-1-yl)piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-

5 yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0284]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-

10 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-(piperidin-1-yl)piperidine (22.0 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-(piperidin-1-yl)piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (44.9 mg).

MS (FAB<sup>+</sup>) m/z: 608 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>34</sub>H<sub>47</sub>FN<sub>5</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 608.3612; found, 608.3583.

[0285]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[4-(piperidin-1-yl)piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-(piperidin-1-yl)piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (44.9 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[4-(piperidin-1-yl)piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.5 mg).

MS (FAB<sup>+</sup>) m/z: 474 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>41</sub>FN<sub>5</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 474.3244; found, 474.3234.

[Example 57]

15 [0286]



[0287]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-methoxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 51, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.0 mg) and 4-




methoxyphenylaniline (22.9 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-methoxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (19.7 mg).

5 MS (FAB<sup>+</sup>) m/z: 429 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 429.2302; found, 429.2330.

[Example 58]

[0288]



10

[0289]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

15

(2S,4S)-1-[[N-(4-Carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.0 mg), along with 1-hydroxybenzotriazole, was dissolved in N,N-dimethylformamide (1.0 mL). While the solution was chilled in an ice bath, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (53.4 mg) was added and the mixture was allowed to warm to room temperature and was stirred for 1 hour. Subsequently, 4-trifluoromethylaniline (23.0  $\mu$ L) was added and the mixture was stirred for additional 12 hours, followed by

20

addition of dimethylaminopyridine (11.3 mg) and stirring for additional 24 hours. The solvent was then evaporated under reduced pressure and the resulting residue was purified by preparative thin-layer chromatography (solvent:


5 dichloromethane: methanol = 4:1) to give (2S,4S)-4-fluoro-1-  
[[N-[4-[N-(4-  
trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile (10.1 mg).

MS (FAB<sup>+</sup>) m/z: 467 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>F<sub>4</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 467.2070; found,  
467.2051.

[Example 59]

[0290]



15 [0291]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-  
adamantylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0292]

20 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-  
adamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and adamantanamine (17.1 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-adamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (38.2 mg). MS (FAB<sup>+</sup>) m/z: 591 (MH<sup>+</sup>). Rf 0.30 (ethyl acetate: hexane = 4:1).

[0293]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-adamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-adamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.2 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-adamantylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (17.4 mg).

MS (FAB<sup>+</sup>) m/z: 457 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 457.2979; found, 457.2990.

[Example 60]

[0294]



[0295]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-1-  
5 ethylpropylamino)carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0296]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-  
10 ethylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-  
4-fluoropyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-Benzyloxycarbonyl-N-[4-  
carboxybicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (40.0 mg), along with 1-  
15 hydroxybenzotriazole, was dissolved in N,N-dimethylformamide  
(0.8 mL). While the solution was chilled in an ice bath, 1-(3-  
dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (41.9  
mg) was added and the mixture was allowed to warm to room  
temperature and was stirred for 2 hours. Subsequently, 1-  
20 ethylpropylamine (13.2  $\mu$ L) was added and the mixture was  
stirred for additional 17 hours. The solvent was evaporated  
under reduced pressure and the resulting residue was dissolved  
in dichloromethane. The organic layer was washed sequentially

with 0.1 N aqueous hydrochloric acid, saturated aqueous sodium bicarbonate solution and saturated brine. The organic layer was then dried over sodium sulfate and was concentrated under reduced pressure. The residue was purified by column

5 chromatography (eluant: ethyl acetate) to give (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-ethylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (46.0 mg).

MS (FAB<sup>+</sup>) m/z: 527 (MH<sup>+</sup>).

10 Rf 0.33 (ethyl acetate).

[0297]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-1-ethylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


15 In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(N-1-ethylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (46.0 mg) was used to  
20 obtain (2S,4S)-4-fluoro-1-[[N-[4-(N-1-ethylpropylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (22.5 mg).

MS (FAB<sup>+</sup>) m/z: 393 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 393.2666; found,  
25 393.2670.

[Example 61]

[0298]



[0299]

- 5 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[(2R)-N-(2-heptyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile
- (2S,4S)-1-[[N-(4-Carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg),
- 10 1-hydroxybenzotriazole (23.7 mg), JANDAJEL-1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (289 mg) and N,N-dimethylformamide (1 mL) were mixed together and the mixture was stirred at room temperature for 3 hours.
- Subsequently, (2R)-2-aminoheptane (46.6 µL) was added and the
- 15 mixture was stirred at room temperature for 17 hours and 40 minutes. This was followed by addition of dichloromethane (0.5 mL) and (2R)-2-aminoheptane (11.6 µL), stirring at room temperature for 4.5 hours, addition of JANDAJEL-1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (96.6 mg) and
- 20 additional stirring at room temperature for 17 hours.
- Subsequently, (isocyanatomethyl)polystyrene (232 mg) was added and the mixture was stirred at room temperature for 2 hours. The insoluble material was filtered and the filtrate was

concentrated under reduced pressure. The resulting residue was then purified by silica gel column (eluant: ethyl acetate: methanol = 10:1) to give (2S,4S)-4-fluoro-1-[[N-[4-[(2R)-N-(2-heptyl)amino]carbonylbicyclo[2.2.2]oct-1-


5 yl]amino]acetyl]pyrrolidine-2-carbonitrile (9.0 mg).

MS (FAB<sup>+</sup>) m/z: 421 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 421.2979; found, 421.2983.

[Example 62]

10 [0300]



[0301]

Synthesis of (2S,4S)-1-[[N-[4-[4-[(3R)-3-(N,N-diethylcarbamoyl)piperidin-1-yl]piperidin-1-

15 yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0302]

Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-[(3R)-3-

20 N,N-diethylcarbamoylpiperidin-1-yl]piperidin-1-

yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-

benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (74.9 mg) and (3R)-N,N-diethyl-1-(piperidin-4-yl)piperidine-3-carboxamide (61.0 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-[(3R)-3-(N,N-diethylcarbamoyl)piperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (71.7 mg).

MS (FAB<sup>+</sup>) m/z: 707 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>39</sub>H<sub>56</sub>N<sub>6</sub>O<sub>5</sub> (MH<sup>+</sup>): calcd, 707.4296; found, 707.4294.

[0303]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-[4-[(3R)-3-(N,N-diethylcarbamoyl)piperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-4-fluoro-1-[[N-benzyloxycarbonyl-N-[4-[4-[(3R)-3-(N,N-diethylcarbamoyl)piperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (66.7 mg) was used to obtain (2S,4S)-1-[[N-[4-[4-[(3R)-3-(N,N-diethylcarbamoyl)piperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.4 mg).




MS (FAB<sup>+</sup>) m/z: 573 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>31</sub>H<sub>50</sub>FN<sub>6</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 573.3928; found, 573.3905.

[Example 63]

5 [0304]



[0305]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[(2R)-N-(2-octyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

10 (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) was suspended in dichloromethane (1 mL). To the suspension, trichloroacetonitrile (31.0  $\mu$ L) and triphenylphosphine (81.1 mg) in dichloromethane (0.5 mL) were added and the mixture was stirred at room temperature for 2 hours. Subsequently, (piperidinomethyl)polystyrene (150 mg) and (2R)-2-aminooctane (57.1  $\mu$ L) were sequentially added at 0°C and the mixture was stirred at room temperature for 21 hours. This was followed by addition of (isocyanatomethyl)polystyrene (232 mg), stirring at room temperature for 1 hour, addition of water (3 mL) and dichloromethane (2 mL), and further stirring at room temperature for 50 minutes. The reaction mixture was then


loaded onto an Isolute HM-N column and was extracted 5 times with 2ml dichloromethane. The dichloromethane extracts were combined and concentrated under reduced pressure. The resulting residue was purified by silica gel column (eluant: 5 dichloromethane: methanol = 50:1) to give (2S,4S)-4-fluoro-1-[[N-[4-[(2R)-N-(2-octyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (30.3 mg).

MS (FAB<sup>+</sup>) m/z: 435 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>40</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 435.3135; found, 10 435.3103.

[Example 64]

[0306]



[0307]

15 Synthesis of (2S,4S)-1-[[N-[4-[4-[(3R)-3-ethoxycarbonylpiperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0308]

20 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-[(3R)-3-ethoxycarbonylpiperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 4, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (101 mg) and ethyl (3R)-1-(piperidin-4-yl)piperidine-3-carboxylate (84.3mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-[(3R)-3-ethoxycarbonylpiperidin-1-yl]piperidine-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (115 mg).

MS (FAB<sup>+</sup>) m/z: 680 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>37</sub>H<sub>51</sub>FN<sub>5</sub>O<sub>6</sub> (MH<sup>+</sup>): calcd, 680.3823; found, 680.3824.

[0309]

Step 2:

Synthesis of (2S,4S)-1-[[N-[4-[4-[(3R)-3-ethoxycarbonylpiperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[4-[(3R)-3-ethoxycarbonylpiperidin-1-yl]piperidin-1-yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (110 mg) was used to obtain (2S,4S)-1-[[N-[4-[4-[(3R)-3-ethoxycarbonylpiperidin-1-yl]piperidin-1-


yl]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (36.0 mg).

MS (FAB<sup>+</sup>) m/z: 546 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>45</sub>FN<sub>5</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 546.3456; found,  
5 546.3452.

[Example 65]

[0310]



[0311]

10 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-2-octyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (2S)-amino-octane (57.1  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-2-octyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (13.4 mg).

MS (FAB<sup>+</sup>) m/z: 435 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>40</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 435.3135; found, 435.3163.

[Example 66]

[0312]



[0313]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(1R)-1-phenyl-1-ethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (1R)-1-phenylethylamine (43.4  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(1R)-1-phenyl-1-ethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile (22.8 mg).

MS (FAB<sup>+</sup>) m/z: 427 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 427.2509; found, 427.2511.

[Example 67]

[0314]



[0315]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-methylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-4-methylthiazole (38.8 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-methylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (11.5 mg).

MS (FAB<sup>+</sup>) m/z: 420 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>27</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 420.1870; found, 420.1837.

[Example 68]

[0316]



15 [0317]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-2-heptyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (2S)-aminoheptane (51.1  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-2-heptyl]amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (22.5 mg).

MS (FAB<sup>+</sup>) m/z: 421 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 421.2979; found, 421.2983.

5 [Example 69]

[0318]



[0319]

10 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-3,3-dimethyl-2-butyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (2S)-3,3-dimethyl-2-butylamine (41.4  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-3,3-dimethyl-2-butyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (20.5 mg).

MS (FAB<sup>+</sup>) m/z: 407 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 407.2822; found, 407.2809.

[Example 70]

[0320]



[0321]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-phenylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-phenylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (32.7mg) were obtained.

MS (FAB<sup>+</sup>) m/z: 482 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>29</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 482.2026; found, 482.2018.

[Example 71]

[0322]



[0323]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[(tetrahydropyran-4-



yl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl)amino]acetyl]pyrrolidine-2-carbonitrile  
[0324]

Step 1:

5 Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-  
[(tetrahydropyran-4-yl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 13, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-(benzotriazol-1-yl)oxycarbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-aminotetrahydrofuran hydrochloride (15.5 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[(tetrahydropyran-4-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (46.4 mg).

MS (FAB<sup>+</sup>) m/z: 541 (MH<sup>+</sup>).

[0325]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[(tetrahydropyran-4-yl)amino]carbonylbicyclo[2.2.2]oct-1-  
20 yl)amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[(tetrahydropyran-4-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (44.4 mg) was used to obtain


(2S,4S)-4-fluoro-1-[[N-[(tetrahydropyran-4-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (13.5 mg).

MS (FAB<sup>+</sup>) m/z: 407 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 407.2458; found, 407.2410.

[Example 72]

[0326]



10 [0327]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(1S)-1-phenyl-1-ethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0328]

15 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-[(1S)-1-phenyl-1-ethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-Benzyloxycarbonyl-N-(4-

20 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (40.0 mg) and 1-

hydroxybenzotriazole (20.1 mg) were dissolved in N,N-

dimethylformamide (0.8 mL). While the solution was chilled in

an ice bath, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (41.9 mg) was added and the mixture was stirred at room temperature for 2 hours. Subsequently, (1S)-1-phenylethylamine (14.5  $\mu$ L) was added and the mixture was further stirred at room temperature for 16.5 hours. The reaction mixture was concentrated under reduced pressure and the resulting residue was dissolved in dichloromethane (2 mL). The dichloromethane solution was washed sequentially with 0.1 mol/L hydrochloric acid, saturated aqueous sodium bicarbonate solution and saturated brine. The solution then was dried over anhydrous sodium sulfate and was concentrated under reduced pressure. The resulting residue was purified by a silica gel column (eluant: ethyl acetate) to give ((2S,4S)-1-[[N-benzyloxycarbonyl-N-[4-[N-[(1S)-1-phenyl-1-ethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (51.5 mg).

MS (FAB<sup>+</sup>) m/z: 561 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>32</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 561.2877; found, 561.2860.

[0329]

Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(1S)-1-phenyl-1-ethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 5, (2S,4S)-1-[[N-

benzyloxycarbonyl-N-(4-[(1S)-N-(1-phenyl-1-ethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (49.0 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(1S)-1-phenyl-1-


5 ethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (24.2 mg).

MS (FAB<sup>+</sup>) m/z: 427 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 427.2509; found, 427.2502.

10 [Example 73]

[0330]



[0331]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-

15 chlorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-chloroaniline (43.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-chlorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (25.6 mg).

MS (FAB<sup>+</sup>) m/z: 433 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>27</sub>ClFN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 433.1807; found, 433.1816.

[Example 74]

[0332]



[0333]

Synthesis of (2S,4S)-1-[[N-(4-[N-(4-acetylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-aminoacetophenone (46.0 mg) were used to obtain (2S,4S)-1-[[N-(4-[N-(4-acetylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (24.4 mg).

MS (FAB<sup>+</sup>) m/z: 441 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 441.2302; found, 441.2291.

[Example 75]

[0334]



[0335]


Synthesis of (2S,4S)-1-[[N-[4-[N-(benzathiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-aminobenzothiazole (51.1 mg) was used to obtain (2S,4S)-1-[[N-[4-[N-(benzathiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (28.0 mg).  
MS (FAB<sup>+</sup>) m/z: 456 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 456.1870; found, 456.1881.

[Example 76]

[0336]



[0337]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-

ethoxycarbonylmethylthiazol-2-  
yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
5 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethy 2-  
aminothiazole-4-acetate (63.3 mg) were used to obtain (2S,4S)-  
1-[[N-[4-[N-(4-ethoxycarbonylmethylthiazol-2-  
yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
10 fluoropyrrolidine-2-carbonitrile (11.2 mg).

MS (FAB<sup>+</sup>) m/z: 492 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>31</sub>FN<sub>5</sub>O<sub>4</sub>S(MH<sup>+</sup>): calcd, 492.2081; found,  
492.2104.

[Example 77]

15 [0338]



[0339]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-  
methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
20 yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-


fluoropyrrolidine-2-carbonitrile (50.0 mg) and p-toluidine (36.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.2 mg).

5 MS (FAB<sup>+</sup>) m/z: 413 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 413.2353; found, 413.2378.

[Example 78]

[0340]



10

[0341]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-methylsulfonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

15

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-methylsulfonylaniline hydrochloride (71.0 mg) were used to obtain ((2S,4S)-4-fluoro-1-[[N-[4-[N-(4-

20

methylsulfonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (13.8 mg).


MS (FAB<sup>+</sup>) m/z: 477 (MH<sup>+</sup>).



HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>4</sub>S(MH<sup>+</sup>): calcd, 477.1972; found, 477.1984.

[Example 79]

[0342]



[0343]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-carbamoylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-aminobenzamide (46.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-carbamoylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (13.2 mg).

MS (FAB<sup>+</sup>) m/z: 442 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>29</sub>FN<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 442.2254; found, 442.2268.

[Example 80]

[0344]



[0345]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-  
cyanophenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 5 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-  
 aminobenzonitrile (40.0 mg) were used to obtain (2S,4S)-1-[[N-  
 10 [4-[N-(4-cyanophenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (17.6 mg).

MS (FAB<sup>+</sup>) m/z: 424 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>FN<sub>5</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 424.2149; found,  
 424.2129.

15 [Example 81]

[0346]



[0347]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-

trifluoromethylsulfonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-trifluoromethylsulfonylaniline (77.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-trifluoromethylsulfonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (10.8 mg).

MS (FAB<sup>+</sup>) m/z: 531 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>F<sub>4</sub>N<sub>4</sub>O<sub>4</sub>S(MH<sup>+</sup>): calcd, 531.1689; found, 531.1682.

[Example 82]

[0348]



[0349]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-benzoylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-

aminobenzophenone (67.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-benzoylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (11.5 mg).  
MS (FAB<sup>+</sup>) m/z: 503 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 503.2458; found, 503.2439.

[Example 83]

[0350]



10 [0351]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-aminosulfonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and sulfanilamide (59.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-aminosulfonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (18.2 mg).

20 MS (FAB<sup>+</sup>) m/z: 478 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>29</sub>FN<sub>5</sub>O<sub>4</sub>S (MH<sup>+</sup>): calcd, 478.1924; found, 478.1940.

[Example 84]

[0352]



[0353]

5 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-aminobenzotrifluoride (55.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (28.4 mg).

15 MS (FAB<sup>+</sup>) m/z: 467 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>F<sub>4</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 467.2070; found, 467.2083.

[Example 85]

[0354]



20

[0355]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-fluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-fluoroaniline (38.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-fluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.2 mg).

MS (FAB<sup>+</sup>) m/z: 417 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>27</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 417.2102; found, 417.2151.

[Example 86]

[0356]



[0357]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-fluoro-3-trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 5-amino-2-fluorobenzotrifluoride (41.0 mg) were used to obtain (2S,4S)-


4-fluoro-1-[[N-[4-[N-(4-fluoro-3-trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (17.8 mg).

MS (FAB<sup>+</sup>) m/z: 485 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>26</sub>F<sub>5</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 485.1976; found, 485.1945.

[Example 87]

[0358]



10 [0359]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(6-fluorobenzotriazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-(4-Carboxybicyclo[2.2.2]oct-1-

15 yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg),

benzotriazol-1-yl-oxytris(dimethylamino)phosphonium

hexafluorophosphate (137 mg) and dichloromethane (1.5 mL) were

mixed together. While the mixture was maintained at 0°C,

triethylamine (43.1 µL) was added and the mixture was stirred

20 at room temperature for 75 minutes. Subsequently, 2-amino-6-

fluorobenzotriazole (57.2 mg) was added and the mixture was

stirred at room temperature for one day. The resulting mixture

was washed sequentially with water and saturated aqueous

sodium bicarbonate solution. The mixture was then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by a silica gel column (eluant: dichloromethane: methanol = 10:1) to give


(2S,4S)-4-fluoro-1-[[N-[4-[N-(6-fluorobenzothiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (49.7 mg) as a pale yellow solid.

MS (FAB<sup>+</sup>) m/z: 474 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>26</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 474.1775; found, 474.1793.

[Example 88]

[0360]



[0361]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-cyclopropylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-4-cyclopropylthiazole (47.7 mg) were used to obtain (2S,4S)-1-




[ [N-[4-[N-(4-cyclopropylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (43.1 mg).

MS (FAB<sup>+</sup>) m/z: 446 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>29</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 446.2026; found, 446.2017.

[Example 89]

[0362]



10 [0363]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-phenyl-5-ethoxycarbonylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethyl 2-amino-5-phenylthiazole-6-carboxylate (84.5 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-phenyl-5-ethoxycarbonylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (34.1 mg).


MS (FAB<sup>+</sup>) m/z: 554 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>33</sub>FN<sub>5</sub>O<sub>4</sub>S(MH<sup>+</sup>): calcd, 554.2237; found,

554.2235.

[Example 90]

[0364]



5 [0365]

Synthesis of (2S,4S)-1-[[N-[4-[N-(2-chloro-4-trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (80.0 mg) and 4-amino-3-chlorobenzotrifluoride (111 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(2-chloro-4-trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (15.3 mg).

MS (FAB<sup>+</sup>) m/z: 501 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>26</sub>ClF<sub>4</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 501.1680; found, 501.1713.

[Example 91]

20 [0366]



[0367]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-fluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (80.0 mg) and 3-fluoroaniline (63.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-fluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (38.1 mg).

MS (FAB<sup>+</sup>) m/z: 417 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>27</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 417.2102; found, 417.2144.

[Example 92]

[0368]



[0369]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-

chlorophenylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-(4-Carboxybicyclo[2.2.2]oct-1-  
yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg),

5 1-hydroxybenzotriazole (28.0 mg), PS-carbodiimide (240 mg) and  
dichloromethane (4 mL) were mixed together and the mixture was  
stirred at room temperature for 15 minutes. 4-

chlorobenzylamine (19.0  $\mu$ L) was then added and the mixture was  
stirred at room temperature for further 24 hours. Subsequently,

10 MP-carbonate (270 mg) was added and the mixture was stirred at  
room temperature for 5 hours and was left overnight. The

insoluble material in the mixture was filtered and the  
filtrate was concentrated under reduced pressure. The

resulting residue was then purified by a silica gel column

15 (eluant: dichloromethane: methanol = 10:1) to give (2S,4S)-1-  
[[N-[4-[N-(4-

chlorophenylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (29.1 mg).

MS (FAB<sup>+</sup>) m/z: 447 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>29</sub>ClFN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 447.1963; found,  
447.1977.

[Example 93]

[0370]



[0371]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-fluoro-2-  
trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 5 yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-  
 fluorobenzotrifluoride (64.0 mg) were used to obtain (2S,4S)-  
 10 4-fluoro-1-[[N-[4-[N-(4-fluoro-2-  
 trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.8 mg).

MS (FAB<sup>+</sup>) m/z: 485 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>26</sub>F<sub>5</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 485.1976; found,  
 15 485.2004.

[Example 94]

[0372]



[0373]

Synthesis of (2S,4S)-1-[[N-[4-[N-[(1-ethoxycarbonyl-1-methoxyiminomethyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

5 In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethyl 2-amino- $\alpha$ -methoxyiminothiazole-4-acetate (78.0 mg) were used to obtain  
 10 (2S,4S)-1-[[N-[4-[N-[(1-ethoxycarbonyl-1-methoxyiminomethyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (43.4 mg).


MS (FAB<sup>+</sup>) m/z: 535 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>32</sub>FN<sub>6</sub>O<sub>5</sub>S(MH<sup>+</sup>): calcd, 535.2139; found,

15 535.2119

[Example 95]

[0374]



[0375]

20 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-methylphenylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-

carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-methylbenzylamine (41.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-(4-[N-(4-


5 methylphenylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile (18.1 mg).

MS (FAB<sup>+</sup>) m/z: 427 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 427.2509; found, 427.2534.

10 [Example 96]

[0376]



[0377]

Synthesis of (2S,4S)-4-fluoro-1-[[N-(4-[N-(4-

15 trifluoromethylphenylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-

20 (trifluoromethyl)benzylamine (60.0 mg) were used to obtain


(2S,4S)-4-fluoro-1-[[N-(4-[N-(4-trifluoromethylphenylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile (22.0 mg).

MS (FAB<sup>+</sup>) m/z: 481 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>29</sub>F<sub>4</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 481.2227; found, 481.2228.

[Example 97]

5 [0378]



[0379]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(1-methylethyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

10

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-isopropylaniline (46.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(1-methylethyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (36.3 mg).

15

MS (FAB<sup>+</sup>) m/z: 441 (MH<sup>+</sup>).


HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 441.2666; found, 441.2687.

20

[Example 98]

[0380]





[0381]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(2-  
hydroxyethyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-  
 5 yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-(4-  
 aminophenyl)ethanol (47.0 mg) were used to obtain (2S,4S)-4-  
 10 fluoro-1-[[N-[4-[N-[4-(2-  
 hydroxyethyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-  
 yl]amino]acetyl]pyrrolidine-2-carbonitrile (35.6 mg).

MS (FAB<sup>+</sup>) m/z: 443 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 443.2548; found,  
 15 443.2452.

[Example 99]

[0382]



[0383]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-  
butylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-butylaniline (51.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-butylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (28.2 mg). MS (FAB<sup>+</sup>) m/z: 455 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 455.2822; found, 455.2859.

[Example 100]

[0384]



[0385]

15 Synthesis of (2S,4S)-1-[[N-[4-[N-(4-ethoxycarbonylmethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethyl 4-aminophenylacetate (61.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-ethoxycarbonylmethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (33.6 mg).

MS (FAB<sup>+</sup>) m/z: 485 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 485.2564; found, 485.2576.

5 [Example 101]

[0386]



[0387]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-morpholinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-morpholinylaniline (61.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-morpholinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (33.0 mg).

MS (FAB<sup>+</sup>) m/z: 484 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>35</sub>FN<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 484.2724; found, 484.2726.

[Example 102]

[0388]



[0389]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-

5 phenylmethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-

10 aminodiphenylmethane (62.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-


phenylmethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (31.1 mg).

MS (FAB<sup>+</sup>) m/z: 489 (MH<sup>+</sup>).

15 HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 489.2666; found, 489.2638.

[Example 103]

[0390]



20 [0391]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(1,1-dimethylethyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-tert-butylaniline (51.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(1,1-dimethylethyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (21.1 mg).

MS (FAB<sup>+</sup>) m/z: 455 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 455.2822; found, 455.2821.

[Example 104]

[0392]



[0393]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-ethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-


fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-ethylaniline (40.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-ethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (24.7 mg).

5 MS (FAB<sup>+</sup>) m/z: 427 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 427.2509; found, 427.2469.

[Example 105]

[0394]



10

[0395]

Synthesis of (2S,4S)-1-[[N-[4-[N-(3-chlorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


15 In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 3-chloroaniline (36.0  $\mu$ L) were used to obtain (2S,4S)-1-[[N-[4-[N-(3-chlorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (23.3 mg).

20

MS (FAB<sup>+</sup>) m/z: 433 (MH<sup>+</sup>).


HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>27</sub>ClFN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 433.1807; found,

[0396]



Synthesis of (2S,4S)-1-[[N-[4-[N-(2-chlorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[ 0398 ]



[0399]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and m-toluidine (25.8  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (18.3 mg).

MS (FAB<sup>+</sup>) m/z: 413 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 413.2353; found, 413.2367.

[Example 108]

[0400]



[0401]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(1-



methylpropyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-


5 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-sec-butylaniline (51.5 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-(1-methylpropyl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (13.0 mg).

10 MS (FAB<sup>+</sup>) m/z: 455 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 455.2822; found, 455.2829.

[Example 109]

[0402]



15

[0403]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-octylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

20 In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-octylaniline


(70.5 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-octylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.8 mg).

MS (FAB<sup>+</sup>) m/z: 511 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>30</sub>H<sub>44</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 511.3448; found, 511.3455.

[Example 110]

[0404]



10 [0405]

Synthesis of (2S,4S)-1-[[N-[4-[N-(2,4-dichlorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2,4-dichloroaniline (55.1 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(2,4-dichlorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (25.5 mg).

20 MS (FAB<sup>+</sup>) m/z: 467 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>26</sub>Cl<sub>2</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 467.1417; found, 467.1441.

[Example 111]

[0406]



[0407]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(2-

5 pyridyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-4-(2-pyridyl)thiazole (60.3 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(2-pyridyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (11.6 mg).

MS (FAB<sup>+</sup>) m/z: 483 (MH<sup>+</sup>).

15 HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>28</sub>FN<sub>6</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 483.1978; found, 483.1966.

[Example 112]

[0408]



[0409]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(4-pyridyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


5 In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-4-(4-pyridyl)thiazole (60.3 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(4-pyridyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.2 mg).

MS (FAB<sup>+</sup>) m/z: 483 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>28</sub>FN<sub>6</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 483.1978; found, 483.2014.

15 [Example 113]

[0410]



[0411]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-


carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and o-toluidine (36.3  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (22.6 mg).

MS (FAB<sup>+</sup>) m/z: 413 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 413.2353; found, 413.2384.

[Example 114]

10 [0412]



[0413]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2,4-dimethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

15


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2,4-dimethylaniline (42.3  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(2,4-dimethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (22.4 mg).

MS (FAB<sup>+</sup>) m/z: 427 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 427.2509; found, 427.2490.

[Example 115]

[0414]



5

[0415]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-cyclohexylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

10 In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-cyclohexylaniline (60.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-cyclohexylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (22.3 mg).


15

MS (FAB<sup>+</sup>) m/z: 481 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 481.2979; found, 481.2932.

[Example 116]

20 [0416]



[0417]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-  
pentylloxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 5 yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-  
 pentylloxyaniline (61.0 mg) were used to obtain (2S,4S)-4-  
 10 fluoro-1-[[N-[4-[N-(4-  
 pentylloxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 yl]amino]acetyl]pyrrolidine-2-carbonitrile (37.6 mg).

MS (FAB<sup>+</sup>) m/z: 485 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 485.2928; found,  
 15 485.2905.

[Example 117]

[0418]



[0419]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-styrylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-aminostilbene (66.4 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-styrylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (21.3 mg).

MS (FAB<sup>+</sup>) m/z: 501 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>30</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 501.2666; found, 501.2637.

[Example 118]

[0420]



[0421]

Synthesis of (2S,4S)-1-[[N-[4-[N-(2-chloro-4-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-chloro-4-




methylaniline (41.8  $\mu$ L) were used to obtain (2S,4S)-1-[[N-[4-[N-(2-chloro-4-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (22.6 mg).

MS (FAB<sup>+</sup>) m/z: 447 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>29</sub>ClFN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 447.1963; found, 447.2000.

[Example 119]

[0422]



10 [0423]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-trifluoromethoxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.3 mg) and 4-trifluoromethoxyaniline (60.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-trifluoromethoxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (28.2 mg).


MS (FAB<sup>+</sup>) m/z: 483 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>F<sub>4</sub>N<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 483.2019; found,

483.1989.

[Example 120]

[0424]



5 [0425]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-fluoro-4-morpholinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.3 mg) and 3-fluoro-4-morpholinylaniline (62.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-fluoro-4-morpholinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (28.6 mg).

MS (FAB<sup>+</sup>) m/z: 502 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>34</sub>F<sub>2</sub>N<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 502.2630; found, 502.2647.

[Example 121]

20 [0426]



[0427]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-ethynylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, ((2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-ethynylaniline (40.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-ethynylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (10.0 mg).

MS (FAB<sup>+</sup>) m/z: 423 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 423.2196; found, 423.2204.

[Example 122]

[0428]



[0429]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-thienylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 92, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-

5 thienylmethylamine (38.5 mg) were used to obtain (2S,4S)-4-

fluoro-1-[[N-[4-[N-(2-

thienylmethyl)amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (48.8 mg).

MS (FAB<sup>+</sup>) m/z: 419 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 419.1917; found, 419.1937.

[Example 123]

[0430]



15 [0431]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-piperidinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

20 fluoropyrrolidine-2-carbonitrile (50.0 mg) and N-(4-

aminophenyl)piperidine (61.4 mg) were used to obtain (2S,4S)-


4-fluoro-1-[[N-[4-[N-(4-piperidinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.6 mg).

MS (FAB<sup>+</sup>) m/z: 482 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>37</sub>FN<sub>5</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 482.2931; found, 482.2913.

[Example 124]

[0432]



10 [0433]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-vinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-aminostyrene (46.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-vinylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (13.1 mg).

20 MS (FAB<sup>+</sup>) m/z: 425 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 425.2353; found, 425.2314.

[Example 125]

[0434]



[0435]

- 5 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2,4,6-trimethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (75.0 mg) and 2,4,6-trimethylaniline (71.6  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(2,4,6-trimethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (29.1 mg).

15 MS (FAB<sup>+</sup>) m/z: 441 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 441.2666; found, 441.2659.

[Example 126]

[0436]



[0437]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-chloro-2-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


5 In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-chlorotoluene (48.2 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-chloro-2-methylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (16.8 mg).

MS (FAB<sup>+</sup>) m/z: 447 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>29</sub>ClFN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 447.1963; found, 447.1973.

[Example 127]

15 [0438]



[0439]

Synthesis of (2S,4S)-1-[[N-[4-[N-[4-(4-chlorophenyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

20

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-

carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-4-(4-chlorophenyl)thiazole (71.7 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-4-(4-chlorophenyl)thiazol-2-


5 yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (23.3 mg).

MS (FAB<sup>+</sup>) m/z: 516 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>28</sub>ClFN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 516.1636; found, 516.1620.

10 [Example 128]

[0440]



[0441]

Synthesis of (2S,4S)-1-[[N-[4-[N-(1,3,4-thiadiazol-2-

15 yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-1,3,4-thiadiazole (34.4 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (20.9 mg).


MS (FAB<sup>+</sup>) m/z: 407 (MH<sup>+</sup>).



HRMS (FAB<sup>+</sup>) for C<sub>18</sub>H<sub>24</sub>FN<sub>6</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 407.1665; found, 407.1620.

[Example 129]

[0442]



5

[0443]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(2,2-dimethylethyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


10 In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-4-tert-butylthiazole (53.1 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(2,2-dimethylethyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (35.9 mg).

MS (FAB<sup>+</sup>) m/z: 462 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>33</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 462.2339; found, 462.2286.

20 [Example 130]

[0444]



[0445]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(5-methyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-methyl-1,3,4-thiadiazole (39.2 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(5-methyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (40.4 mg).

MS (FAB<sup>+</sup>) m/z: 421 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>26</sub>FN<sub>6</sub>O<sub>2</sub>S (MH<sup>+</sup>): calcd, 421.1822; found, 421.1862.

[Example 131]

[0446]



[0447]

Synthesis of (2S,4S)-1-[[N-[4-[N-(5-ethyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-ethyl-1,3,4-thiadiazole (43.9 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(5-ethyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (34.8 mg).

MS (FAB<sup>+</sup>) m/z: 435 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>28</sub>FN<sub>6</sub>O<sub>2</sub>S (MH<sup>+</sup>): calcd, 435.1978; found, 435.1990.

[Example 132]

[0448]



15 [0449]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4,4-difluorocyclohexyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

20 [0450]

Step 1:

Synthesis of 4-benzyloxycarbonylamino-N-(4,4-

difluorocyclohexyl)bicyclo[2.2.2]octane-1-carboxamide

1-Hydroxybenzotriazole (138mg), 4-

benzyloxycarbonylbicyclo[2.2.2]octane-1-carboxylic acid (72.0 mg) and N,N-dimethylformamide (8 mL) were mixed together.

5 While the mixture was chilled in an ice bath, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (265 mg) was added and the mixture was stirred at room temperature for 1 hour. Subsequently, a mixture of 4,4-difluorocyclohexylamine hydrochloride (108 mg), triethylamine  
10 (105  $\mu$ L) and N,N-dimethylformamide (2 mL) was added and the resultant mixture was stirred at room temperature for 18 hours and was then concentrated under reduced pressure. Water (10 mL) was added to the residue and the solution was extracted with ethyl acetate (3  $\times$  10 mL). The ethyl acetate extracts  
15 were combined, washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by a silica gel column (eluant: hexane: ethyl acetate = 3:1) to give 4-benzyloxycarbonylamino-N-(4,4-  
20 difluorocyclohexyl)bicyclo[2.2.2]octane-1-carboxamide (57.0 mg) as a white solid.

MS (EI) m/z: 420 ( $M^+$ ).

[0451]

Step 2:

25 Synthesis of 4-amino-N-(4,4-

difluorocyclohexyl)bicyclo[2.2.2]octane-1-carboxamide

4-Benzyloxycarbonylamino-N-(4,4-difluorocyclohexyl)bicyclo[2.2.2]octane-1-carboxamide (55.4 mg) was dissolved in tetrahydrofuran (6 mL). To this solution, 10% palladium carbon (20.0 mg) was added and the mixture was stirred at room temperature for 6 hours in a hydrogen atmosphere. The reaction mixture was filtered to remove the catalyst and the filtrate was concentrated under reduced pressure. The resulting residue was purified by a silica gel column (eluant: ethyl acetate: methanol = 10:1) to give 4-amino-N-(4,4-difluorocyclohexyl)bicyclo[2.2.2]octane-1-carboxamide (38.1 mg) as a white solid.

MS (EI) m/z: 286 (M<sup>+</sup>).

[0452]

Step 3:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4,4-difluorocyclohexyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

4-Amino-N-(4,4-difluorocyclohexyl)bicyclo[2.2.2]octane-1-carboxamide (31.8 mg), potassium carbonate (16.4 mg) and N,N-dimethylformamide (1.5 mL) were mixed together. To the mixture, (2S,4S)-1-bromoacetyl-4-fluoropyrrolidine-2-carbonitrile (26.3 mg) in N,N-dimethylformamide (1 mL) was added dropwise at room temperature and the mixture was stirred for 90 minutes.

Subsequently, the mixture was concentrated under reduced


pressure and the resulting residue was purified by a silica gel column (eluant: dichloromethane: methanol = 10:1) to give (2S,4S)-4-fluoro-1-[[N-[4-[N-(4,4-difluorocyclohexyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (12.0 mg).

MS (FAB<sup>+</sup>) m/z: 441 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>32</sub>F<sub>3</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 441.2477; found, 441.2475.

[Example 133]

10 [0453]



[0454]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-chlorophenyl)-N-methylamino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

15

[0455]

Step 1:

Synthesis of 4-tert-butoxycarbonylamino-N-(4-chlorophenyl)-N-methylbicyclo[2.2.2]octane-1-carboxamide

20 4-tert-Butoxycarbonylaminobicyclo[2.2.2]octane-1-carboxylic acid (101 mg) was dissolved in dichloromethane (2 mL). To this solution, trichloroacetonitrile (74.0  $\mu$ L) and

triphenylphosphine (196 mg) in dichloromethane (1.5 mL) were sequentially added and the mixture was stirred at room temperature for 2 hours. This was followed by addition of triethylamine (0.18 mL) and 4-chloro-N-methylaniline (98.6 µL) and stirring at room temperature for additional 5.5 hours. The reaction mixture was then poured into aqueous citric acid (5 mL) and was extracted with ethyl acetate (3 x 10 mL). The ethyl acetate extracts were combined, washed with saturated aqueous sodium bicarbonate solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by a silica gel column (eluant: hexane: ethyl acetate = 2:1) to give 4-tert-butoxycarbonylamino-N-(4-chlorophenyl)-N-methylbicyclo[2.2.2]octane-1-carboxamide (91.4 mg) as a white powder.

[0456]

Step 2:

Synthesis of 4-amino-N-(4-chlorophenyl)-N-methylbicyclo[2.2.2]octane-1-carboxamide

4-tert-Butoxycarbonylamino-N-(4-chlorophenyl)-N-methylbicyclo[2.2.2]octane-1-carboxamide (80.0 mg) was mixed with a 4mol/L dioxane solution of hydrogen chloride (1.2 mL) and the mixture was stirred at room temperature for 40 minutes. The crystallized product was collected by filtration and was suspended in water (0.8 mL). While the suspension was chilled

in an ice bath, a 1mol/L aqueous solution of sodium hydroxide (0.3 mL) was added and the mixture was extracted with dichloromethane (4 x 3 mL). The dichloromethane extracts were combined, washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. This gave 4-amino-N-(4-chlorophenyl)-N-methylbicyclo[2.2.2]octane-1-carboxamide (39.8 mg) as a white solid.

[0457]

Step 3:

10 Synthesis of (2S,4S)-1-[[N-[4-[N-(4-chlorophenyl)-N-methylamino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 132, 4-amino-N-(4-chlorophenyl)-N-methylbicyclo[2.2.2]octane-1-carboxamide (30.0 mg) and (2S,4S)-4-fluoropyrrolidine-2-carbonitrile (24.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-chlorophenyl)-N-methylamino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (30.4 mg).


MS (FAB<sup>+</sup>) m/z: 447 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>29</sub>ClFN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 447.1963; found, 447.1994.

[Example 134]

[0458]





[0459]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-  
acetamidophenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 5 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4'-  
 aminoacetanilide (51.0 mg) were used to obtain (2S,4S)-1-[[N-  
 10 [4-[N-(4-acetamidophenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (11.9 mg).

MS (FAB<sup>+</sup>) m/z: 456 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>31</sub>FN<sub>5</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 456.2411; found,  
 456.2403.

15 [Example 135]

[0460]



[0461]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2R)-1-hydroxy-2-

butyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-


5 fluoropyrrolidine-2-carbonitrile (50.0 mg) and (1R)-2-amino-1-butanol (30.1 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2R)-1-hydroxy-2-butyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (11.3 mg).

MS (FAB<sup>+</sup>) m/z: 395 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 395.2458; found, 395.2420.

[Example 136]

[0462]



15 [0463]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(5-trifluoromethyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

20 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-trifluoromethyl-1,3,4-thiadiazole (57.5 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(5-trifluoromethyl-1,3,4-

thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (12.0 mg).


MS (FAB<sup>+</sup>) m/z: 475 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>23</sub>F<sub>4</sub>N<sub>6</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 475.1539; found,

5 475.1557.

[Example 137]

[0464]



[0465]

10 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(4-fluorophenyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-4-(4-fluorophenyl)thiazole (66.1 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(4-fluorophenyl)thiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (50.6 mg).

20 MS (FAB<sup>+</sup>) m/z: 500 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>28</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 500.1932; found, 500.1978.

[Example 138]

[0466]



[0467]

5 Synthesis of (2S,4S)-1-[[N-[4-[N-(5-cyclopropyl-1,3,4-  
thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-cyclopropyl-1,3,4-thiathiazole (48.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(5-cyclopropyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (42.2 mg).

15 MS (FAB<sup>+</sup>) m/z: 447 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>28</sub>FN<sub>6</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 447.1978; found, 447.2007.

[Example 139]

[0468]



[0469]

Synthesis of (2S,4S)-1-[[N-[4-[N-[4-(benzothiazol-2-yl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


5 In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-(benzothiazol-2-yl)aniline (51.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-[4-(benzothiazol-2-yl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (10.4 mg).

MS (FAB<sup>+</sup>) m/z: 532 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>31</sub>FN<sub>5</sub>O<sub>3</sub>S(MH<sup>+</sup>): calcd, 532.2183; found, 532.2158.

15 [Example 140]

[0470]



[0471]

Synthesis of (2S,4S)-1-[[N-[4-[N-[4-(pyrrolidin-1-yl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-


carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-(pyrrolidin-1-yl)aniline (50.2 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-[4-(pyrrolidin-1-yl)phenyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (13.7 mg).

MS (FAB<sup>+</sup>) m/z: 468 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>35</sub>FN<sub>5</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 468.2775; found, 468.2738.

10 [Example 141]

[0472]



[0473]

15 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-methyl-4-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0474]

Step 1:

20 Synthesis of 4-tert-butoxycarbonylamino-N-(2-methyl-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 133, 4-tert-butoxycarbonylaminobicyclo[2.2.2]octane-1-carboxylic acid (150 mg) and 2-methyl-4-trifluoromethylaniline (2152  $\mu$ L) were used

to obtain 4-tert-butoxycarbonylamino-N-(2-methyl-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (92.5 mg).

MS (FAB<sup>+</sup>) m/z: 427 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>30</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 427.2209; found, 427.2237.

[0475]

Step 2:

Synthesis of 4-amino-N-(2-methyl-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide  
10

In a similar manner to Example 133, 4-tert-butoxycarbonylamino-N-(2-methyl-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (84.6 mg) was used to obtain 4-amino-N-(2-methyl-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (59.8 mg).

MS (FAB<sup>+</sup>) m/z: 327 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>17</sub>H<sub>22</sub>F<sub>3</sub>N<sub>2</sub>O (MH<sup>+</sup>): calcd, 327.1684; found, 327.1711.

20 [0476]

Step 3:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-methyl-4-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

25 In a similar manner to Example 132, 4-amino-N-(2-methyl-

4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide  
(59.8 mg) and (2S,4S)-1-bromoacetyl-4-fluoropyrrolidine-2-  
carbonitrile (43.1 mg) were used to obtain (2S,4S)-1-[[N-[4-  
[N-(2-methyl-4-


5 trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (48.4 mg).

MS (FAB<sup>+</sup>) m/z: 481 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>29</sub>F<sub>4</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 481.2227; found,  
481.2247.

10 [Example 142]

[0477]



[0478]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-chloro-2-

15 trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0479]

Step 1:

Synthesis of 4-tert-butoxycarbonylamino-N-(4-chloro-2-

20 trifluoromethyl-phenyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 133, 4-tert-  
butoxycarbonylaminobicyclo[2.2.2]octane-1-carboxylic acid (150



mg) and 2-amino-5-chlorobenzotrifluoride (173  $\mu$ L) were used to obtain 4-tert-butoxycarbonylamino-N-(4-chloro-2-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (79.7 mg).

5 MS (FAB<sup>+</sup>) m/z: 447 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>27</sub>ClF<sub>3</sub>N<sub>2</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 447.1662; found, 447.1631.

[0480]

Step 2:

10 Synthesis of 4-amino-N-(4-chloro-2-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 133, 4-tert-butoxycarbonylamino-N-(4-chloro-2-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (76.9 mg) was used to obtain 4-amino-N-(2-methyl-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (43.0 mg).

MS (FAB<sup>+</sup>) m/z: 347 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>16</sub>H<sub>19</sub>ClF<sub>3</sub>N<sub>2</sub>O (MH<sup>+</sup>): calcd, 347.1138; found, 347.1172.

[0481]

Step 3:


25 Synthesis of (2S,4S)-1-[[N-[4-[N-(4-chloro-2-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 132, 4-amino-N-(4-chloro-2-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (38.2 mg) and (2S,4S)-1-bromoacetyl-4-fluoropyrrolidine-2-carbonitrile (24.2 mg) were used to obtain (2S,4S)-1-[[N-[4-  
5 [N-(4-chloro-2-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (21.9 mg).  
MS (FAB<sup>+</sup>) m/z: 501 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>26</sub>ClF<sub>4</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 501.1680; found,  
10 501.1662.

[Example 143]

[0482]



[0483]

15 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-propylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-propylaniline (46.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-propylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (19.1 mg).

MS (FAB<sup>+</sup>) m/z: 441 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 441.2666; found, 441.2672.

[Example 144]

5 [0484]



[0485]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-  
pentylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-

10 yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-pentylaniline (55.5 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-pentylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (15.7 mg).

MS (FAB<sup>+</sup>) m/z: 469 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 469.2979; found, 469.2977.

20 [Example 145]

[0486]



[0487]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-fluoro-4-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0488]

Step 1:

Synthesis of 4-tert-butoxycarbonylamino-N-(2-fluoro-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 133, 4-tert-butoxycarbonylamino-N-(2-fluoro-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxylic acid (100 mg) and 4-amino-3-fluorobenzotrifluoride (106  $\mu$ L) were used to obtain 4-tert-butoxycarbonylamino-N-(2-fluoro-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (58.6 mg).

MS (FAB<sup>+</sup>) m/z: 431 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>27</sub>F<sub>4</sub>N<sub>2</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 431.1958; found, 431.1970.

[0489]

Step 2:

Synthesis of 4-amino-N-(2-fluoro-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 133, 4-tert-butoxycarbonylamino-N-(2-fluoro-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (55.0 mg) was used to obtain 4-amino-N-(2-fluoro-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (36.2 mg).

[0490]

Step 3:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-fluoro-4-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 133, 4-amino-N-(2-fluoro-4-trifluoromethylphenyl)bicyclo[2.2.2]octane-1-carboxamide (33.0 mg) and (2S,4S)-1-bromoacetyl-4-fluoropyrrolidine-2-carbonitrile (23.5 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-fluoro-4-trifluoromethylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (19.6 mg).

MS (FAB<sup>+</sup>) m/z: 485 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>26</sub>F<sub>5</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 485.1976; found, 485.1983.

[Example 146]

[0491]



[0492]

Synthesis of (2S,4S)-4-fluoro-1-[[N-(4-[N-[(2S)-1-fluoro-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

(2S,4S)-1-[[N-(4-Carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (100 mg) and 1-hydroxybenzotriazole (61.5 mg) were dissolved in dichloromethane (4 mL). While this solution was chilled in an ice bath, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (119 mg) was added and the mixture was stirred at room temperature for 1 hour. Subsequently, a mixture of (2S)-1-fluoro-2-propylamine hydrochloride (32.0 mg), triethylamine (56.0  $\mu$ L) and dichloromethane (2 mL) was added and the resulting mixture was stirred at room temperature for further 8 hours. Following addition of water, the dichloromethane layer was collected. The dichloromethane layer was washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by a silica gel column (eluant: dichloromethane: methanol = 10:1) to give (2S,4S)-4-fluoro-1-[[N-(4-[N-[(2S)-1-fluoro-2-


propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile (34.0 mg) as a white powder.

MS (FAB<sup>+</sup>) m/z: 383 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>29</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 383.2259; found, 383.2227.

[Example 147]

[0493]



10 [0494]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-1-methoxy-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile

[0495]

15 Step 1:

Synthesis of (2S,4S)-1-[[N-benzyloxycarbonyl-N-[(2S)-1-methoxy-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 72, (2S,4S)-1-[[N-benzyloxycarbonyl-N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (40.0 mg) and (2S)-1-methoxy-2-propylamine (10.2 mg) were used to obtain (2S,4S)-1-[[N-benzyloxycarbonyl-N-[(2S)-1-methoxy-2-

20

propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (47.5 mg).

MS (FAB<sup>+</sup>) m/z: 529 (MH<sup>+</sup>).

[0496]

5 Step 2:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-1-methoxy-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


10 In a similar manner to Example 5, (2S,4S)-1-[[N-benzyloxycarbonyl-N-[(2S)-1-methoxy-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (34.3 mg) was used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-1-methoxy-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (11.2 mg).

MS (FAB<sup>+</sup>) m/z: 395 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 395.2458; found, 395.2426.

[Example 148]

20 [0497]



[0498]

Synthesis of (2S,4S)-1-[[N-[4-[N-(5-adamantyl-1,3,4-



thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-


5 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 5-adamantyl-2-amino-1,3,4-thiadiazole (80.0 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(5-adamantyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (56.7 mg).

10 MS (FAB<sup>+</sup>) m/z: 541 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>28</sub>H<sub>38</sub>FN<sub>6</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 541.2761; found, 541.2782.

[Example 149]

[0499]



15

[0500]

Synthesis of (2S,4S)-1-[[N-[4-[N-(5-ethoxycarbonylmethylthio)-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

20 In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethyl [(5-amino-1,3,4-thiadiazol-2-yl)thio]acetate (74.6 mg) were used


to obtain (2S,4S)-1-[[N-[4-[N-(5-ethoxycarbonylmethylthio-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (16.4 mg).

MS (FAB<sup>+</sup>) m/z: 525 (MH<sup>+</sup>).

- 5 HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>30</sub>FN<sub>6</sub>O<sub>4</sub>S<sub>2</sub> (MH<sup>+</sup>): calcd, 525.1754; found, 525.1771.

[Example 150]

[0501]



- 10 [0502]

Synthesis of (2S,4S)-1-[[N-[4-(N,N-dibutylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0503]

- 15 Step 1:

Synthesis of 4-benzyloxycarbonylamino-N,N-dibutylbicyclo[2.2.2]octane-1-carboxamide

- Dibutylamine (94.4  $\mu$ L) and triethylamine (77.9  $\mu$ L) were dissolved in dichloromethane (2 mL). While this solution was chilled in a salt/ice bath, 4-benzyloxycarbonylbicyclo[2.2.2]octane-1-carbonyl chloride (150 mg) in dichloromethane (2 mL) was added dropwise and the mixture was stirred for 40 minutes and was concentrated under
- 20

reduced pressure. Ethyl acetate (30 mL) was then added to the resulting residue and the mixture was washed sequentially with water (1.5 mL), a 2 mol/L aqueous sodium hydroxide solution (1.5 mL), water (1.5 mL) and saturated brine (1.5 mL). The mixture was then dried over anhydrous sodium sulfate and concentrated under reduced pressure. Purification of the resulting residue by a silica gel column (eluant: hexane: ethyl acetate = 3:1) gave 4-benzyloxycarbonylamino-N,N-dibutylbicyclo[2.2.2]octane-1-carboxamide (171 mg) as a white solid.

MS (FAB<sup>+</sup>) m/z: 415 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>39</sub>N<sub>2</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 415.2961; found, 415.2987.

[0504]

Step 2:

Synthesis of 4-amino-N,N-dibutylbicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 132, 4-benzyloxycarbonylamino-N,N-dibutylbicyclo[2.2.2]octane-1-carboxamide (159 mg) was used to obtain 4-amino-N,N-dibutylbicyclo[2.2.2]octane-1-carboxamide (107 mg).

MS (FAB<sup>+</sup>) m/z: 281 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>17</sub>H<sub>33</sub>N<sub>2</sub>O (MH<sup>+</sup>): calcd, 281.2593; found, 281.2624.

[0505]

Step 3:

Synthesis of (2S,4S)-1-[[N-[4-(N,N-  
dibutylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 87, 4-amino-N,N-  
5 dibutylbicyclo[2.2.2]octane-1-carboxamide (58.5 mg) and  
(2S,4S)-4-fluoropyrrolidine-2-carbonitrile (49.0 mg) were used  
to obtain (2S,4S)-1-bromoacetyl-1-[[N-[4-(N,N-  
dibutylamino)carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (18.9 mg).

10 MS (FAB<sup>+</sup>) m/z: 435 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>40</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 435.3135; found,  
435.3156.

[Example 151]

[0506]



15

[0507]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(1,1-  
dimethylethyloxycarbonyl)phenyl]amino]carbonylbicyclo[2.2.2]oc  
t-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

20 In a similar manner to Example 63, (2S,4S)-1-[[N-(4-  
carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
fluoropyrrolidine-2-carbonitrile (120 mg) and 1,1-


dimethylethyl 4-aminobenzoate (158 mg) were used to obtain  
 (2S,4S)-4-fluoro-1-[[N-[4-[N-[4-(1,1-  
 dimethylethyloxycarbonyl)phenyl]amino]carbonylbicyclo[2.2.2]oc  
 t-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (20.1 mg).

5 MS (FAB<sup>+</sup>) m/z: 499 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 499.2721; found,  
 499.2721.

[Example 152]

[0508]



10

[0509]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-  
 carboxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-  
 yl]amino]acetyl]pyrrolidine-2-carbonitrile trifluoroacetate

15

(2S,4S)-4-Fluoro-1-[[N-[4-[N-[4-(1,1-  
 dimethylethyloxycarbonyl)phenyl]amino]carbonylbicyclo[2.2.2]oc  
 t-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (23.0mg) was  
 dissolved in dichloromethane (0.2 mL). To this solution,  
 trifluoroacetic acid (0.2 mL) was added and the mixture was  
 20 stirred at room temperature for 1 hour. Subsequently, the  
 reaction mixture was concentrated under reduced pressure to  
 give (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-


carboxyphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile trifluoroacetate (24.5 mg).

MS (FAB<sup>+</sup>) m/z: 443 (MH<sup>+</sup>).

5 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>28</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 443.2095; found, 443.2077.

[Example 153]

[0510]



10 [0511]

Synthesis of (2S,4S)-4-fluoro-1-[[N-(4-[[N-[(2R)-1-fluoro-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 146, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (100 mg) and (2R)-1-fluoro-2-propylamine hydrochloride (32.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-(4-[[N-[(2R)-1-fluoro-2-propyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]pyrrolidine-2-carbonitrile (41.9 mg).

20 MS (FAB<sup>+</sup>) m/z: 383 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>29</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 383.2259; found, 383.2229.

[Example 154]

[0512]



[0513]

5 Synthesis of (2S,4S)-1-[[N-[4-[N-(5-ethoxycarbonyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethyl (5-amino-1,3,4-thiadiazole-2-carboxylate) (58.9 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(5-ethoxycarbonyl-1,3,4-thiadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (31.9 mg).

15 MS (FAB<sup>+</sup>) m/z: 479 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>28</sub>FN<sub>6</sub>O<sub>4</sub>S(MH<sup>+</sup>): calcd, 479.1877; found, 479.1916.

[Example 155]

[0514]



20

[0515]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-adamantylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-adamantyl-2-aminothiazole (79.7 mg) were used to obtain (2S,4S)-1-[[N-[4-[N-(4-adamantylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (29.5 mg).

MS (FAB<sup>+</sup>) m/z: 540 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>29</sub>H<sub>39</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 540.2809; found, 540.2816.

[Example 156]

[0516]



[0517]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[5-[2-(4-methylphenylsulfonyl)ethyl]-1,3,4-thiadiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-[4-



methylphenylsulfonyl)ethyl]-1,3,4-thiadiazole (96.4 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[5-[2-(4-methylphenylsulfonyl)ethyl]-1,3,4-thiadiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-


5 yl]amino]acetyl]pyrrolidine-2-carbonitrile (30.6 mg).

MS (FAB<sup>+</sup>) m/z: 589 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>27</sub>H<sub>34</sub>FN<sub>6</sub>O<sub>4</sub>S<sub>2</sub> (MH<sup>+</sup>): calcd, 589.2067; found, 589.2081.

[Example 157]

10 [0518]



[0519]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(5-methyl-1,3,4-oxadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-

15 yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-methyl-1,3,4-oxadiazole (33.7 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(5-methyl-1,3,4-oxadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (14.5 mg).


20 4-fluoro-1-[[N-[4-[N-(5-methyl-1,3,4-oxadiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-

MS (FAB<sup>+</sup>) m/z: 405 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>26</sub>FN<sub>6</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 405.2050; found, 405.2075.

[Example 158]

[0520]



[0521]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[5-(4-morpholinyl)-1,3,4-oxadiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-(2-amino-1,3,4-oxadiazole-5-yl)morpholine (57.9 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[5-(4-morpholinyl)-1,3,4-oxadiazol-2-yl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (10.1 mg).

MS (FAB<sup>+</sup>) m/z: 476 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>31</sub>FN<sub>7</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 476.2422; found, 476.2456.

[Example 159]

[0522]



[0523]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-4-hydroxy-2-butyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 146, ((2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (100 mg) and (2S)-3-aminobutanol hydrochloride (39.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(2S)-4-hydroxy-2-butyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (20.8 mg).

MS (FAB<sup>+</sup>) m/z: 395 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 395.2458; found, 395.2462.

[Example 160]

[0524]



[0525]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-methylbutyl)amino]carbonylbicyclo[2.2.2]oct-1-

yl)amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and isoamylamine

5 (39.5  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(3-methylbutyl)amino]carbonylbicyclo[2.2.2]oct-1-

yl)amino]acetyl]pyrrolidine-2-carbonitrile (22.3 mg).


MS (FAB<sup>+</sup>) m/z: 393 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>21</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 393.2666; found,

10 393.2679.

[Example 161]

[0526]



[0527]

15 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(3,3-  
dimethylbutyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl)amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-[4-carboxybicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

20 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 3,3-

dimethylbutylamine (45.8  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(3,3-

dimethylbutyl)amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (24.6 mg).

MS (FAB<sup>+</sup>) m/z: 407 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>36</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 407.2822; found, 407.2779.

5 [Example 162]

[0528]



[0529]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-

10 methoxyethyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-

15 methoxyethylamine (29.6  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(2-

methoxyethyl)amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (31.2 mg).

MS (FAB<sup>+</sup>) m/z: 381 (MH<sup>+</sup>).

20 HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>30</sub>FN<sub>4</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 381.2302; found, 381.2306.

[Example 163]

[0530]



[0531]

Synthesis of (2S,4S)-1-[[N-[4-(N-  
cyclopentylmethylamino)carbonylbicyclo[2.2.2]oct-1-  
 5 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 87, (2S,4S)-1-[[N-(4-  
 carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-  
 fluoropyrrolidine-2-carbonitrile (70.0 mg) and  
 cyclopentylmethylamine hydrochloride (72.2 mg) were used to  
 10 obtain (2S,4S)-1-[[N-[4-(N-  
 cyclopentylmethylamino)carbonylbicyclo[2.2.2]oct-1-  
 yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (19.4 mg).

MS (FAB<sup>+</sup>) m/z: 405 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>34</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 405.2666; found,  
 15 405.2698.

[Example 164]

[0532]



[0533]

20 Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-(N-(4-

heptyl)amino]carbonylbicyclo[2.2.2]oct-1-  
yl)amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-


5 fluoropyrrolidine-2-carbonitrile (50.0 mg) and 4-heptylamine (50.9  $\mu$ L) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(4-heptyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (36.4 mg).

MS (FAB<sup>+</sup>) m/z: 421 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 421.2979; found, 421.2968.

[Example 165]

[0534]



15 [0535]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-((1R,2S)-2-fluoro-1-  
cyclopropyl]amino]carbonylbicyclo[2.2.2]oct-1-  
yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0536]

20 Step 1:

Synthesis of 4-benzyloxycarbonylamino-N-((1R,2S)-2-fluoro-1-  
cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 150, 4-

benzyloxycarbonylbicyclo[2.2.2]octane-1-carbonyl chloride (200 mg) and (1R,2S)-2-fluoro-1-cyclopropylamine p-toluene sulfonate (184 mg) were used to obtain 4-benzyloxycarbonylamino-N-[(1R,2S)-2-fluoro-1-cyclopropyl]bicyclo[2.2.2]octane-1-carboxamide (189 mg).

[0537]

Step 2:

Synthesis of 4-amino-N-[(1R,2S)-2-fluoro-1-cyclopropyl]bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 132, 4-benzyloxycarbonylamino-N-[(1R,2S)-2-fluoro-1-cyclopropyl]bicyclo[2.2.2]octane-1-carboxamide (189 mg) was used to obtain 4-amino-N-[(1R,2S)-2-fluoro-1-cyclopropyl]bicyclo[2.2.2]octane-1-carboxamide (59.1 mg).

[0538]

Step 3:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-[(1R,2S)-2-fluoro-1-cyclopropyl]amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 87, 4-amino-N-[(1R,2S)-2-fluoro-1-cyclopropyl]bicyclo[2.2.2]octane-1-carboxamide (47.5 mg) and (2S,4S)-1-bromoacetyl-4-fluoropyrrolidine-2-carbonitrile (49.3 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-[(1R,2S)-2-fluoro-1-cyclopropyl]amino]carbonylbicyclo[2.2.2]oct-1-




yl]amino]acetyl]pyrrolidine-2-carbonitrile (34.6 mg).

MS (FAB<sup>+</sup>) m/z: 381 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>19</sub>H<sub>27</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 381.2102; found, 381.2128.

5 [Example 166]

[0539]



[0540]

Synthesis of (2S,4S)-1-[[N-[4-[N-(1-ethoxycarbonyl-1-

10 cyclopropyl)amino]carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

[0541]

Step 1:

Synthesis of 4-benzyloxycarbonylamino-N-(1-ethoxycarbonyl-1-

15 cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 150, 4-benzyloxycarbonylbicyclo[2.2.2]octane-1-carbonyl chloride (200 mg) and ethyl 1-amino-1-cyclopropylcarboxylate hydrochloride (123 mg) were used to obtain 4-benzyloxycarbonylamino-N-  
20 [(1R,2S)-1-ethoxycarbonyl-1-cyclopropyl]bicyclo[2.2.2]octane-1-carboxamide (217 mg).

[0542]

Step 2:

Synthesis of 4-amino-N-(1-ethoxycarbonyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 132, 4-benzyloxycarbonylamino-N-[(1R,2S)-1-ethoxycarbonyl-1-cyclopropyl]bicyclo[2.2.2]octane-1-carboxamide (205 mg) was  
5 used to obtain 4-amino-N-(1-ethoxycarbonyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide (124 mg).

MS (FAB<sup>+</sup>) m/z: 281 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>15</sub>H<sub>25</sub>N<sub>2</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 281.1865; found,  
10 281.1856.

[0543]

Step 3:

Synthesis of (2S,4S)-1-[[N-[4-[N-(1-ethoxycarbonyl-1-cyclopropyl)amino]carbonyl]bicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile  
15


In a similar manner to Example 87, 4-amino-N-(1-ethoxycarbonyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide (62.6 mg) and (2S,4S)-1-bromoacetyl-4-fluoropyrrolidine-2-carbonitrile (52.5 mg) were used to obtain  
20 (2S,4S)-1-[[N-[4-[N-(1-ethoxycarbonyl-1-cyclopropyl)amino]carbonyl]bicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (20.2 mg).

MS (FAB<sup>+</sup>) m/z: 435 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>22</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 435.2408; found,  
25 435.2408.

[Example 167]

[0544]



5 [0545]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(1-phenyl-1-cyclopropyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

[0546]

10 Step 1:

Synthesis of 4-benzyloxycarbonylamino-N-(1-phenyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 150, 4-benzyloxycarbonylbicyclo[2.2.2]octane-1-carbonyl chloride (480  
15 mg) and 1-phenyl-1-cyclopropylamine (99.3 mg) were used to obtain 4-benzyloxycarbonylamino-N-(1-phenyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide (255 mg).

MS (FAB<sup>+</sup>) m/z: 419 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>26</sub>H<sub>31</sub>N<sub>2</sub>O<sub>3</sub> (MH<sup>+</sup>): calcd, 419.2335; found,  
20 419.2345.

[0547]

Step 2:

Synthesis of 4-amino-N-(1-phenyl-1-

cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide

In a similar manner to Example 132, 4-benzyloxycarbonylamino-N-(1-phenyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide (255 mg) was  
5 used to obtain 4-amino-N-(1-phenyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide (118 mg).

MS (FAB<sup>+</sup>) m/z: 285 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>18</sub>H<sub>25</sub>N<sub>2</sub>O (MH<sup>+</sup>): calcd, 285.1967; found, 285.1982.

[0548]

10 Step 3:

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[N-(1-phenyl-1-cyclopropyl)amino]carbonyl]bicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile


In a similar manner to Example 87, 4-amino-N-(1-phenyl-1-cyclopropyl)bicyclo[2.2.2]octane-1-carboxamide (50.0 mg) and  
15 (2S,4S)-1-bromoacetyl-4-fluoropyrrolidine-2-carbonitrile (41.9 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-[4-[N-(1-phenyl-1-cyclopropyl)amino]carbonyl]bicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (38.1 mg).

20 MS (FAB<sup>+</sup>) m/z: 439 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 439.2509; found, 439.2512.

[Example 168]

[0549]



[0550]

Synthesis of (2S,4S)-1-[[N-(4-[[N-[(1R)-1-cyclohexylethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile


In a similar manner to Example 63, (2S,4S)-1-[[N-[4-carboxybicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (50.0 mg) and (1R)-1-cyclohexylethylamine (49.7  $\mu$ L) were used to obtain (2S,4S)-1-[[N-(4-[[N-[(1R)-1-cyclohexylethyl]amino]carbonylbicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (29.9 mg).

MS (FAB<sup>+</sup>) m/z: 433 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>24</sub>H<sub>38</sub>FN<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 433.2979; found, 433.2996.

[Example 169]

[0551]



[0552]

Synthesis of (2S,4S)-4-fluoro-1-[[N-[4-[[N-(3-methylthiazol-2-yl)amino]carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-[4-carboxybicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

fluoropyrrolidine-2-carbonitrile (50.0 mg) and 2-amino-5-

5 methylthiazole (38.8 mg) were used to obtain (2S,4S)-4-fluoro-

1-[[N-[4-[N-(3-methylthiazol-2-

yl]amino]carbonylbicyclo[2.2.2]oct-1-


yl]amino]acetyl]pyrrolidine-2-carbonitrile (9.5 mg).

MS (FAB<sup>+</sup>) m/z: 420 (MH<sup>+</sup>).

10 HRMS (FAB<sup>+</sup>) for C<sub>20</sub>H<sub>27</sub>FN<sub>5</sub>O<sub>2</sub>S(MH<sup>+</sup>): calcd, 420.1870; found, 420.1874.

[Example 170]

[0553]



15 [0554]

Synthesis of (2S,4S)-1-[[N-[4-[N-(4-

ethoxycarbonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-

yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-

20 fluoropyrrolidine-2-carbonitrile (50.0 mg) and ethyl 4-

aminobenzoate (56.0 mg) were used to obtain (2S,4S)-1-[[N-[4-

[N-(4-ethoxycarbonylphenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (24.9 mg).


MS (FAB<sup>+</sup>) m/z: 471 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>25</sub>H<sub>32</sub>FN<sub>4</sub>O<sub>4</sub> (MH<sup>+</sup>): calcd, 471.2408; found,

5 471.2412.

[Example 171]

[0555]



[0556]

10 Synthesis of (2S,4S)-4-fluoro-1-[[N-(4-[N-(3-trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile

In a similar manner to Example 63, (2S,4S)-1-[[N-(4-carboxybicyclo[2.2.2]oct-1-yl)amino]acetyl]-4-fluoropyrrolidine-2-carbonitrile (80.0 mg) and 3-aminobenzotrifluoride (92.0 mg) were used to obtain (2S,4S)-4-fluoro-1-[[N-(4-[N-(3-trifluorophenyl)amino]carbonylbicyclo[2.2.2]oct-1-yl]amino]acetyl]pyrrolidine-2-carbonitrile (23.2 mg).

20 MS (FAB<sup>+</sup>) m/z: 467 (MH<sup>+</sup>).

HRMS (FAB<sup>+</sup>) for C<sub>23</sub>H<sub>27</sub>F<sub>4</sub>N<sub>4</sub>O<sub>2</sub> (MH<sup>+</sup>): calcd, 467.2070; found, 467.2087.

[0557]

<Test Example 1> [Test for the ability of the compounds of the invention to inhibit of dipeptidylpeptidase IV activity]

The concentration of free 7-amino-4-methyl-coumarin (AMC) generated by hydrolysis of H-Gly-Pro-AMC·HBr substrate by plasma dipeptidylpeptidase IV was determined by fluorometry.

#### Method

A 20  $\mu$ L of buffer (25mmol/L hepes, 140mmol/L sodium chloride, 1% bovine serum albumin, 80mmol/L magnesium chloride hexahydrate, pH 7.4) containing each compound was added to 20  $\mu$ L of plasma diluted 8-fold with saline in a well of a 96-well flat bottom plate. The plate was left at room temperature for 5 minutes and 10  $\mu$ L of 0.1mmol/L H-Gly-Pro-AMC·HBr solution was added to each well to initiate the reaction. The plate was left in a dark environment at room temperature for 20 minutes, at which point 20 $\mu$ L 25% acetic acid was added to terminate the reaction. Using a fluorescent plate reader, the free AMC concentration was determined by exciting the samples at 355 nm and measuring the fluorescence intensity at 460 nm. Using Prism 3.02 (GraphPad Software), the results were analyzed to determine the 50% inhibitory concentration (IC<sub>50</sub>). The results are shown in Table 1.

[0558]



**Table 1: *In vitro* dipeptidylpeptidase IV inhibition**

Test compound	I C 5 0 (n mol/L)
Example 1	0 . 8 9
Example 8	0 . 8 3
Example 16	0 . 0 8 2
Example 52	0 . 0 5 7
Compound A	3 . 3

[0559]

Compound A: (2S)-1-[[ (3-hydroxy-1-adamantyl)amino]acetyl]-2-cyanopyrrolidine (LAF-237)

5 [0560]

<Test Example 2> [Test for the inhibition of dipeptidylpeptidase IV activity in mice by oral administration of the compounds of the invention]

Each compound was suspended in 0.3% sodium carboxymethylcellulose to a concentration of 0.1 mg/mL. The preparation was orally administered to 8-week old male ICR mice (Charles River Laboratories Japan) at a dose of 10 mL/kg. Using an EDTA 2K-treated capillary tube, blood samples were collected from the tail vein before administration and 30 minutes after administration. The blood samples were centrifuged at 6000 rpm for 2 minutes to separate plasma. The enzymatic activity was determined using the same procedure as in Test Example 1. The inhibition was determined from the decrease in the enzymatic activity from the initial activity

20 (% inhibition = {(activity before administration - activity

after administration)/(activity before administration)} x 100).

The results are shown in Table 2.

[0561]

Table 2: Inhibition of plasma dipeptidylpeptidase IV activity  
in mice by oral administration

Test compound	% inhibition
Example 1	71
Example 9	87
Example 15	66
Example 30	77
Example 52	70
Compound A	81

5 [0562]

Compound A: (2S)-1-[[[(3-hydroxy-1-adamantyl)amino]acetyl]-2-cyanopyrrolidine (LAF-237)

[0563]

<Test Example 3> [Oral glucose tolerance test in mice]

10           The compound of the present invention of Example 58 was  
suspended in 0.3% sodium carboxymethylcellulose (CMC-Na,  
Sigma). Seven weeks old male ICR mice (Charles River  
Laboratories Japan) were acclimatized for 1 week. During the  
acclimatization period, the animals were allowed to freely  
15 consume standard feed (CE-2, Clea Japan) and water. The ICR  
mice reaching 8-weeks old were fasted for 16 hours.  
Subsequently, the animals were orally administered 0.3%CMC-Na  
(10 mL/kg) or Compound 1 (1 mg/kg, 10 mL/kg) and were  
immediately administered a glucose solution orally at a dose  
20 of 5 g/kg. Using an EDTA 2K-treated capillary tube, blood  
samples were collected from the tail vein before

administration of glucose solution and 15, 30, 60, and 120 minutes after administration. The blood glucose level was determined using glucose B-test Wako (Wako Pure Chemical Industries). The results were shown in means  $\pm$  standard errors.

5 Statistical analysis was performed using t-test with a significant level of less than 5%. The results are shown in Fig. 1.

[0564]

<Test Example 4> [Test for the efficacy of the compounds of  
10 the invention against drug-induced hypoleukocytosis]

The efficacy of the compounds of the present invention against drug-induced hypoleukocytosis was evaluated by conducting an experiment according to the method described by Okabe et al (Japanese Pharmacology and Therapeutics, Vol. 19,  
15 No. 6 (1991): p55).

[0565]

Eight weeks old male ICR mice (Charles River Laboratories Japan) were intraperitoneally administered a single dose of cyclophosphamide (200 mg/kg) on Day 0. Starting from the  
20 following day, control group was given saline and test group was orally administered the compound of the present invention (1 to 200 mg/kg) once or twice a day over a five day period. Blood samples were collected 2, 4, 6, and 8 days after the beginning of the test and the white blood cell count was  
25 monitored over time. The white blood cell count of the test

group at a given time was compared with the white blood cell count before administration of cyclophosphamide to evaluate the efficacy of the compound of the present invention against the drug-induced hypoleukocytosis. The results indicate that the decrease in the white blood cell count is significantly suppressed in the group administered the compound of the present invention as compared to control group.

[0566]

<Test Example 5> [Test for the ability of the compounds of the invention to increase the blood G-CSF level]

Seven weeks old male ICR mice (Charles River Laboratories Japan) were used. Control group was given saline and test group was orally administered the compound of the present invention (1 to 200 mg/kg) once or twice a day over a five day period. Mice were anesthetized on the day following the cessation of administration and blood samples were collected. Plasma G-CSF level was determined using mouse G-CSF ELISA kit (R&D SYSTEM). The results indicate that the plasma G-CSF level was significantly increased in the group administered the compound of the present invention as compared to control group.

INDUSTRIAL APPLICABILITY

[0567]

As set forth, the compounds of the present invention are novel bicycloamide derivatives and pharmaceutically acceptable salts thereof that effectively inhibit DPP-IV. Pharmaceutical

compositions that contain the present compound as an active ingredient are useful in the prevention and/or treatment of diabetes and associated diabetic complications, as well as in the prevention and/or treatment of other diseases that involve

5 DPP-IV.